


*Dissipation from the one-particle-irreducible effective action*

Stefan Flörchinger (Heidelberg U.)

INT Program: Multi-Scale Problems Using Effective Field Theories,  
Seattle, May 8, 2018.



UNIVERSITÄT  
HEIDELBERG  
ZUKUNFT  
SEIT 1386



## *Effective dissipation*

- dissipation is **generation** of entropy
- von Neumann definition

$$S = -\text{Tr}\rho \ln \rho$$

- entropy measures information
  - maximal information for pure state with  $S = 0$
  - minimal information for thermal state  $S = \max. |_{E, \vec{p}, N}$
- unitary evolution **conserves** entropy!
- what information is really accessible and relevant?

## *Entanglement entropy*

- consider splitting of system into two parts  $A + B$
- reduced density matrix

$$\rho_A = \text{Tr}_B\{\rho\}$$

- entanglement entropy

$$S_A = -\text{Tr}_A\{\rho_A \ln \rho_A\}$$

- C-theorem & A-theorem
- local entropy production  $\leftrightarrow$  entanglement generation

## *Dissipation and effective field theory*

- RG equations for the dissipative terms?
- universality in the effective dissipative sector?
- what dissipative terms are relevant for dynamics close to (quantum) phase transitions?

## *Close-to-equilibrium situations*

- out-of-equilibrium situations
- close-to-equilibrium: description by field expectation values and thermodynamic fields
- more complete description by following more fields explicitly
- example: viscous fluid dynamics plus additional fields
- usually discussed in terms of
  - phenomenological constitutive relations
  - as a limit of kinetic theory
  - in AdS/CFT
- want non-perturbative formulation in terms of QFT concepts
- analytic continuation as an alternative to Schwinger-Keldysh
- direct generalization of equilibrium formalism

## Local equilibrium states

- dissipation: energy and momentum get transferred to a heat bath
- even if one starts with pure state  $T = 0$  initially, dissipation will generate nonzero temperature
- close-to-equilibrium situations: dissipation is local
- convenient to use general coordinates with metric

$$g_{\mu\nu}(x)$$

- need approximate **local** equilibrium description with temperature  $T(x)$  and fluid velocity  $u^\mu(x)$ , will appear in combination

$$\beta^\mu(x) = \frac{u^\mu(x)}{T(x)}$$

- **global** thermal equilibrium corresponds to  $\beta^\mu$  Killing vector

$$\nabla_\mu \beta_\nu(x) + \nabla_\nu \beta_\mu(x) = 0$$

## Local equilibrium

- similarity between local density matrix and translation operator

$$e^{\beta^\mu(x) \mathcal{P}_\mu} \longleftrightarrow e^{i\Delta x^\mu \mathcal{P}_\mu}$$

- functional integral with periodicity in imaginary direction


$$\phi(x^\mu - i\beta^\mu(x)) = \pm\phi(x^\mu)$$

- partition function  $Z[J]$ , Schwinger functional  $W[J]$  in Euclidean domain


$$Z[J] = e^{W_E[J]} = \int D\phi e^{-S_E[\phi] + \int_x J\phi}$$

- first defined on **Euclidean manifold**  $\Sigma \times M$  at constant time
- approximate local equilibrium at all times: hypersurface  $\Sigma$  can be shifted

(a) Global thermal equilibrium



(b) Local thermal equilibrium



## *Effective action*

- defined in euclidean domain by Legendre transform

$$\Gamma_E[\Phi] = \int_x J_a(x) \Phi_a(x) - W_E[J]$$

with expectation values

$$\Phi_a(x) = \frac{1}{\sqrt{g(x)}} \frac{\delta}{\delta J_a(x)} W_E[J]$$

- quantum or 1-PI effective action has correlation functions including all quantum fluctuations !
- euclidean field equation

$$\frac{\delta}{\delta \Phi_a(x)} \Gamma_E[\Phi] = \sqrt{g(x)} J_a(x)$$

resembles classical equation of motion for  $J = 0$

- need analytic continuation to obtain a viable equation of motion



## Two-point functions

- consider homogeneous background fields and global equilibrium

$$\beta^\mu = \left( \frac{1}{T}, 0, 0, 0 \right)$$

- propagator and inverse propagator

$$\frac{\delta^2}{\delta J_a(-p)\delta J_b(q)} W_E[J] = G_{ab}(i\omega_n, \mathbf{p}) \delta(p - q)$$

$$\frac{\delta^2}{\delta \Phi_a(-p)\delta \Phi_b(q)} \Gamma_E[\Phi] = P_{ab}(i\omega_n, \mathbf{p}) \delta(p - q)$$

- from definition of effective action

$$\sum_b G_{ab}(p) P_{bc}(p) = \delta_{ac}$$


## Spectral representation

- Källén-Lehmann spectral representation

$$G_{ab}(\omega, \mathbf{p}) = \int_{-\infty}^{\infty} dz \frac{\rho_{ab}(z^2 - \mathbf{p}^2, z)}{z - \omega}$$

with  $\rho_{ab} \in \mathbb{R}$

- correlation functions can be analytically continued in  $\omega = -u^\mu p_\mu$
- branch cut or poles on real frequency axis  $\omega \in \mathbb{R}$  but nowhere else
- different propagators follow by evaluation of  $G_{ab}$  in different regions



$$\Delta_{ab}^M(p) = G_{ab}(i\omega_n, \mathbf{p})$$

$$\Delta_{ab}^R(p) = G_{ab}(p^0 + i\epsilon, \mathbf{p})$$

$$\Delta_{ab}^A(p) = G_{ab}(p^0 - i\epsilon, \mathbf{p})$$

$$\Delta_{ab}^F(p) = G_{ab}(p^0 + i\epsilon \text{ sign}(p^0), \mathbf{p})$$

## Inverse propagator

- spectral representation for  $G_{ab}$  implies that *inverse propagator*  $P_{ab}(\omega, \mathbf{p})$ 
  - can have zero-crossings for  $\omega = p^0 \in \mathbb{R}$
  - has in general branch-cut for  $\omega = p^0 \in \mathbb{R}$
- so far reference frame with  $u^\mu = (1, 0, 0, 0)$
- more general: analytic continuation with respect to

$$\omega = -u^\mu p_\mu$$

- use **decomposition**

$$P_{ab}(p) = P_{1,ab}(p) - i s_1(-u^\mu p_\mu) P_{2,ab}(p)$$

with **sign function**

$$s_1(\omega) = \text{sign}(\text{Im } \omega)$$

- both functions  $P_{1,ab}(p)$  and  $P_{2,ab}(p)$  are regular (no discontinuities)

## Sign operator in position space

[Floerchinger, JHEP 1609 (2016) 099]

- in position space, **sign function** becomes **operator**

$$s_I(-u^\mu p_\mu) = \text{sign}(\text{Im}(-u^\mu p_\mu))$$

$$\rightarrow \text{sign}\left(\text{Im}\left(iu^\mu \frac{\partial}{\partial x^\mu}\right)\right) = \text{sign}\left(\text{Re}\left(u^\mu \frac{\partial}{\partial x^\mu}\right)\right) = s_R\left(u^\mu \frac{\partial}{\partial x^\mu}\right)$$

- geometric representation in terms of Lie derivative

$$s_R(\mathcal{L}_u) \quad \text{or} \quad s_R(\mathcal{L}_\beta)$$

- **sign operator** appears also in analytically continued quantum effective action  $\Gamma[\Phi]$

## *Analytically continued 1 PI effective action*

[Floerchinger, JHEP 1609 (2016) 099]

- analytically continued quantum effective action defined by analytic continuation of correlation functions
- quadratic part

$$\Gamma_2[\Phi] = \frac{1}{2} \int_{x,y} \Phi_a(x) \left[ P_{1,ab}(x-y) + P_{2,ab}(x-y)_{\text{SR}} \left( u^\mu \frac{\partial}{\partial y^\mu} \right) \right] \Phi_b(y)$$

- higher orders correlation functions less understood: no spectral representation
- use inverse Hubbard-Stratonovich trick: terms quadratic in auxiliary field can be integrated out
- allows to understand analytic structures of higher order terms

## *Equations of motion*

- can one obtain **causal** and **real** renormalized equations of motion from the 1 PI effective action?
- naively: time-ordered action / Feynman  $i\epsilon$  prescription:

$$\frac{\delta}{\delta\Phi_a(x)} \Gamma_{\text{time ordered}}[\Phi] = \sqrt{g} J_a(x)$$

- this does not lead to causal and real equations of motion !  
[e.g. Calzetta & Hu: *Non-equilibrium Quantum Field Theory* (2008)]

## Retarded functional derivative

[Floerchinger, JHEP 1609 (2016) 099]

- **real** and **causal dissipative field equations** follow from analytically continued effective action

$$\left. \frac{\delta\Gamma[\Phi]}{\delta\Phi_a(x)} \right|_{\text{ret}} = \sqrt{g}J(x)$$

- to calculate retarded variational derivative determine

$$\delta\Gamma[\Phi]$$

by varying the fields  $\delta\Phi(x)$  including dissipative terms

- set signs according to

$$s_R(u^\mu \partial_\mu) \delta\Phi(x) \rightarrow -\delta\Phi(x), \quad \delta\Phi(x) s_R(u^\mu \partial_\mu) \rightarrow +\delta\Phi(x)$$

- proceed as usual
- opposite choice of sign: field equations for backward time evolution
- leads to causal equations of motion

## Damped harmonic oscillator 1

- equation of motion

$$m\ddot{x} + c\dot{x} + kx = 0$$

or

$$\ddot{x} + 2\zeta\omega_0\dot{x} + \omega_0^2x = 0$$

with  $\omega_0 = \sqrt{k/m}$  and  $\zeta = c/\sqrt{4mk}$

- what is effective action for damped oscillator? This does *not* work:

$$\int \frac{d\omega}{2\pi} \frac{m}{2} x^*(\omega) [\omega^2 + 2i\omega\zeta\omega_0 - \omega_0^2] x(\omega)$$

- consider inverse propagator

$$\omega^2 + 2i s_1(\omega) \omega \zeta \omega_0 - \omega_0^2$$

with

$$s_1(\omega) = \text{sign}(\text{Im } \omega)$$

zero crossings (poles in the eff. propagator) are broadened to branch cut



## Damped harmonic oscillator 2

- take for effective action

$$\begin{aligned}\Gamma[x] &= \int \frac{d\omega}{2\pi} \frac{m}{2} x^*(\omega) [-\omega^2 - 2i s_1(\omega) \omega \zeta \omega_0 + \omega_0^2] x(\omega) \\ &= \int dt \left\{ -\frac{1}{2} m \dot{x}^2 + \frac{1}{2} c x s_R(\partial_t) \dot{x} + \frac{1}{2} k x^2 \right\}\end{aligned}$$

where the second line uses

$$s_1(\omega) = \text{sign}(\text{Im } \omega) \rightarrow \text{sign}(\text{Im } i\partial_t) = \text{sign}(\text{Re } \partial_t) = s_R(\partial_t)$$

- variation gives up to boundary terms

$$\delta\Gamma = \int dt \left\{ m \ddot{x} \delta x + \frac{1}{2} c \delta x s_R(\partial_t) \dot{x} - \frac{1}{2} c \dot{x} s_R(\partial_t) \delta x + k x \delta x \right\}$$

Set now  $s_R(\partial_t) \delta x \rightarrow -\delta x$  and  $\delta x s_R(\partial_t) \rightarrow \delta x$ . Defines  $\frac{\delta\Gamma}{\delta x} \Big|_{\text{ret}}$ .

- equation of motion for forward time evolution

$$\frac{\delta\Gamma}{\delta x} \Big|_{\text{ret}} = m \ddot{x} + c \dot{x} + k x = 0$$

## Scalar field with $O(N)$ symmetry

- consider effective action (with  $\rho = \frac{1}{2}\varphi_j\varphi_j$ )

$$\Gamma[\varphi, g_{\mu\nu}, \beta^\mu] = \int d^d x \sqrt{g} \left\{ \frac{1}{2} Z(\rho, T) g^{\mu\nu} \partial_\mu \varphi_j \partial_\nu \varphi_j + U(\rho, T) + \frac{1}{2} C(\rho, T) [\varphi_j, s_R(u^\mu \partial_\mu)] \beta^\nu \partial_\nu \varphi_j \right\}$$

- variation at fixed metric  $g_{\mu\nu}$  and  $\beta^\mu$  gives

$$\begin{aligned} \delta\Gamma = \int d^d x \sqrt{g} \left\{ Z(\rho, T) g^{\mu\nu} \partial_\mu \delta\varphi_j \partial_\nu \varphi_j + \frac{1}{2} Z'(\rho, T) \varphi_m \delta\varphi_m g^{\mu\nu} \partial_\mu \varphi_j \partial_\nu \varphi_j \right. \\ + U'(\rho, T) \varphi_m \delta\varphi_m \\ + \frac{1}{2} C(\rho, T) [\delta\varphi_j, s_R(u^\mu \partial_\mu)] \beta^\nu \partial_\nu \varphi_j \\ + \frac{1}{2} C(\rho, T) [\varphi_j, s_R(u^\mu \partial_\mu)] \beta^\nu \partial_\nu \delta\varphi_j \\ \left. + \frac{1}{2} C'(\rho, T) \varphi_m \delta\varphi_m [\varphi_j, s_R(u^\mu \partial_\mu)] \beta^\nu \partial_\nu \varphi_j \right\} \end{aligned}$$

- set now  $\delta\varphi_j s_R(u^\mu \partial_\mu) \rightarrow \delta\varphi_j$  and  $s_R(u^\mu \partial_\mu) \delta\varphi_j \rightarrow -\delta\varphi_j$

## *Scalar field with $O(N)$ symmetry*

- field equation becomes

$$-\nabla_{\mu} [Z(\rho, T)\partial^{\mu}\varphi_j] + \frac{1}{2}Z'(\rho, T)\varphi_j\partial_{\mu}\varphi_m\partial^{\mu}\varphi_m \\ + U'(\rho, T)\varphi_j + C(\rho, T)\beta^{\mu}\partial_{\mu}\varphi_j = 0$$

- generalized Klein-Gordon equation with additional damping term

## Causality

[Floerchinger, JHEP 1609 (2016) 099]

- consider derivative of field equation (in flat space with  $\sqrt{g} = 1$ )

$$\left. \frac{\delta}{\delta\Phi_b(y)} \frac{\delta\Gamma}{\delta\Phi_a(x)} \right|_{\text{ret}} = \frac{\delta}{\delta\Phi_b(y)} J_a(x)$$

- inverting this equation gives retarded Green's function

$$\frac{\delta}{\delta J_b(y)} \Phi_a(x) = \Delta_{ab}^R(x, y)$$

- only non-zero for  $x$  future or null to  $y$
- **Causality**: Field expectation value  $\Phi_a(x)$  can only be influenced by the source  $J_b(y)$  in or on the past light cone ✓

## *Where do energy & momentum go?*

- modified variational principle leads to equations of motion with dissipation
- but what happens to the dissipated energy and momentum?
- and other conserved quantum numbers?
- what about entropy production?

## *Energy-momentum tensor expectation value*

- analogous to field equation, obtain by retarded variation

$$\left. \frac{\delta \Gamma[\Phi, g_{\mu\nu}, \beta^\mu]}{\delta g_{\mu\nu}(x)} \right|_{\text{ret}} = -\frac{1}{2} \sqrt{g} \langle T^{\mu\nu}(x) \rangle$$

- leads to Einstein's field equation when  $\Gamma[\Phi, g_{\mu\nu}, \beta^\mu]$  contains Einstein-Hilbert term
- useful to decompose

$$\Gamma[\Phi, g_{\mu\nu}, \beta^\mu] = \Gamma_R[\Phi, g_{\mu\nu}, \beta^\mu] + \Gamma_D[\Phi, g_{\mu\nu}, \beta^\mu]$$

where reduced action  $\Gamma_R$  contains no dissipative / discontinuous terms and  $\Gamma_D$  only dissipative terms

- energy-momentum tensor has two parts

$$\langle T^{\mu\nu} \rangle = (\bar{T}_R)^{\mu\nu} + (\bar{T}_D)^{\mu\nu}$$

## General covariance

- infinitesimal general coordinate transformations as a *gauge transformation* of the metric

$$\delta g_{\mu\nu}^G(x) = g_{\mu\lambda}(x) \frac{\partial \epsilon^\lambda(x)}{\partial x^\nu} + g_{\nu\lambda}(x) \frac{\partial \epsilon^\lambda(x)}{\partial x^\mu} + \frac{\partial g_{\mu\nu}(x)}{\partial x^\lambda} \epsilon^\lambda(x)$$

- temperature / fluid velocity field transforms as vector

$$\delta \beta_G^\mu(x) = -\beta^\nu(x) \frac{\partial \epsilon^\mu(x)}{\partial x^\nu} + \frac{\partial \beta^\mu(x)}{\partial x^\nu} \epsilon^\nu(x)$$

- also fields  $\Phi_a$  transform in some representation, e. g. as scalars

$$\delta \Phi_a^G(x) = \epsilon^\lambda(x) \frac{\partial}{\partial x^\lambda} \Phi_a(x)$$

- reduced action is invariant

$$\Gamma_R[\Phi + \delta \Phi^G, g_{\mu\nu} + \delta g_{\mu\nu}^G, \beta^\mu + \beta_G^\mu] = \Gamma_R[\Phi, g_{\mu\nu}, \beta^\mu]$$

## *Situation without dissipation*

- consider first situation **without dissipation**  $\Gamma[\Phi, g_{\mu\nu}, \beta^\mu] = \Gamma_R[\Phi, g_{\mu\nu}]$
- field equation implies (for  $J = 0$ )

$$\frac{\delta}{\delta\Phi_a(x)} \Gamma_R[\Phi, g_{\mu\nu}] = 0$$

- gauge variation of the metric

$$\delta\Gamma_R = \int d^d x \sqrt{g} \epsilon^\lambda(x) \nabla_\mu \langle T^\mu{}_\lambda(x) \rangle$$

- general covariance  $\delta\Gamma_R = 0$  and field equations imply covariant energy-momentum conservation

$$\nabla_\mu \langle T^\mu{}_\lambda(x) \rangle = 0$$



## Situation with dissipation

[Floerchinger, JHEP 1609 (2016) 099]

- consider now situation **with dissipation**. General covariance of  $\Gamma_R$ :

$$\delta\Gamma_R = \int d^d x \left\{ \frac{\delta\Gamma_R}{\delta\Phi_a} \delta\Phi_a^G + \sqrt{g} \epsilon^\lambda \nabla_\mu (\bar{T}_R)^\mu{}_\lambda + \frac{\delta\Gamma_R}{\delta\beta^\mu} \delta\beta^\mu \right\} = 0$$

- reduced action **not stationary** with respect to field variations

$$\frac{\delta\Gamma_R}{\delta\Phi_a(x)} = - \frac{\delta\Gamma_D}{\delta\Phi_a(x)} \Big|_{\text{ret}} =: -\sqrt{g}(x) M_a(x)$$

- reduced energy-momentum tensor **not conserved**

$$\nabla_\mu (\bar{T}_R)^\mu{}_\lambda(x) = -\nabla_\mu (\bar{T}_D)^\mu{}_\lambda(x)$$

- dependence on  $\beta^\mu(x)$  **cannot be dropped**

$$\frac{\delta\Gamma_R}{\delta\beta^\mu(x)} =: \sqrt{g}(x) K_\mu(x)$$

- general covariance implies **four additional differential equations** that determine  $\beta^\mu$

$$M_a \partial_\lambda \Phi_a + \nabla_\mu (\bar{T}_D)^\mu{}_\lambda = \nabla_\mu [\beta^\mu K_\lambda] + K_\mu \nabla_\lambda \beta^\mu$$

## Entropy production

[Floerchinger, JHEP 1609 (2016) 099]

- contraction of previous equation with  $\beta^\lambda$  gives

$$M_a \beta^\lambda \partial_\lambda \Phi_a + \beta^\lambda \nabla_\mu (\bar{T}_D)^\mu{}_\lambda = \nabla_\mu [\beta^\mu \beta^\lambda K_\lambda]$$

- consider special case

$$\sqrt{g} K_\mu(x) = \frac{\delta \Gamma_R}{\delta \beta^\mu(x)} = \frac{\delta}{\delta \beta^\mu(x)} \int d^d x \sqrt{g} U(T)$$

with grand canonical potential density  $U(T) = -p(T)$  and temperature

$$T = \frac{1}{\sqrt{-g_{\mu\nu} \beta^\mu \beta^\nu}}$$

- using  $s = \partial p / \partial T$  gives entropy current

$$\beta^\mu \beta^\lambda K_\lambda = s^\mu = s u^\mu$$

- local form of **second law of thermodynamics**

$$\nabla_\mu s^\mu = M_a \beta^\lambda \partial_\lambda \Phi_a + \beta^\lambda \nabla_\mu (\bar{T}_D)^\mu{}_\lambda \geq 0$$

## Energy-momentum tensor for scalar field

- analytic action

$$\Gamma[\varphi, g_{\mu\nu}, \beta^\mu] = \int d^d x \sqrt{g} \left\{ \frac{1}{2} Z(\rho, T) g^{\mu\nu} \partial_\mu \varphi_j \partial_\nu \varphi_j + U(\rho, T) \right. \\ \left. + \frac{1}{2} C(\rho, T) [\varphi_j, s_R(u^\mu \partial_\mu)] \beta^\nu \partial_\nu \varphi_j \right\}$$

- energy-momentum tensor

$$\langle T^{\mu\nu}(x) \rangle = Z(\rho, T) \partial^\mu \varphi_j \partial^\nu \varphi_j \\ - \left( g^{\mu\nu} + u^\mu u^\nu T \frac{\partial}{\partial T} \right) \left\{ \frac{1}{2} Z(\rho, T) g^{\mu\nu} \partial_\mu \varphi_j \partial_\nu \varphi_j + U(\rho, T) \right\}$$

- generalizes  $T^{\mu\nu}$  for scalar field and  $T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu + g^{\mu\nu} p$  for ideal fluid with pressure  $p = -U$  and enthalpy density  $\epsilon + p = sT = -T \frac{\partial}{\partial T} U$ .
- general covariance and covariant conservation law imply

$$\nabla_\mu \langle T^{\mu\nu}(x) \rangle = 0 \quad \implies \quad \text{Differential eqs. for } \beta^\mu(x)$$

## Entropy production for scalar field

- entropy current

$$s^\mu = \beta^\mu \beta^\lambda K_\lambda = -\beta^\mu T \frac{\partial}{\partial T} \left\{ \frac{1}{2} Z(\rho, T) g^{\alpha\beta} \partial_\alpha \varphi_j \partial_\beta \varphi_j + U(\rho, T) \right\}$$

- generalized entropy density

$$s_G = -\frac{\partial}{\partial T} \left\{ \frac{1}{2} Z(\rho, T) g^{\alpha\beta} \partial_\alpha \varphi_j \partial_\beta \varphi_j + U(\rho, T) \right\}$$

- entropy generation positive semi-definite for  $C(\rho, T) \geq 0$

$$\nabla_\mu s^\mu = C(\rho, T) (\beta^\mu \partial_\mu \varphi_j) (\beta^\nu \partial_\nu \varphi_j) \geq 0$$

- for fluid at rest  $u^\mu = (1, 0, 0, 0)$

$$\nabla_\mu s^\mu = \dot{s}_G = \frac{C(\rho, T)}{T^2} \dot{\varphi}_j \dot{\varphi}_j$$

entropy increases when  $\varphi_j$  oscillates. Example: reheating after inflation

## *Ideal fluid*

- consider effective action

$$\Gamma[g_{\mu\nu}, \beta^\mu] = \Gamma_R[g_{\mu\nu}, \beta^\mu] = \int d^d x \sqrt{g} U(T)$$

with effective potential  $U(T) = -p(T)$  and temperature

$$T = \frac{1}{\sqrt{-g_{\mu\nu} \beta^\mu \beta^\nu}}$$

- variation of  $g_{\mu\nu}$  at fixed  $\beta^\mu$  lead to ideal fluid form

$$T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu + pg^{\mu\nu}$$

where  $\epsilon + p = Ts = T \frac{\partial}{\partial T} p$  is the enthalpy density

- general covariance or covariant conservation  $\nabla_\mu T^{\mu\nu} = 0$  leads to

$$\begin{aligned} u^\mu \partial_\mu \epsilon + (\epsilon + p) \nabla_\mu u^\mu &= 0, \\ (\epsilon + p) u^\mu \nabla_\mu u^\nu + \Delta^{\nu\mu} \partial_\mu p &= 0. \end{aligned}$$

## Viscous fluid

- analytic action

$$\Gamma[g_{\mu\nu}, \beta^\mu] = \int_x \left\{ U(T) + \frac{1}{4} [g_{\mu\nu}, s_R(\mathcal{L}_u)] (2\eta(T)\sigma^{\mu\nu} + \zeta(T)\Delta^{\mu\nu}\nabla_\rho u^\rho) \right\}$$

with projector

$$\Delta^{\mu\nu} = u^\mu u^\nu + g^{\mu\nu}$$

and

$$\sigma^{\mu\nu} = \left( \frac{1}{2} \Delta^{\mu\alpha} \Delta^{\mu\beta} + \frac{1}{2} \Delta^{\mu\beta} \Delta^{\mu\alpha} - \frac{1}{d-1} \Delta^{\mu\nu} \Delta^{\alpha\beta} \right) \nabla_\alpha u_\beta$$

leads to

$$\langle T^{\mu\nu} \rangle = -\frac{2}{\sqrt{g}} \frac{\delta\Gamma[g_{\mu\nu}, \beta^\mu]}{\delta g_{\mu\nu}} \Big|_{\text{ret}} = (\epsilon + p)u^\mu u^\nu + pg^{\mu\nu} - 2\eta\sigma^{\mu\nu} - \zeta\Delta^{\mu\nu}\nabla_\rho u^\rho$$

- describes viscous fluid with shear viscosity  $\eta(T)$  and bulk viscosity  $\zeta(T)$
- entropy production


$$\nabla_\mu s^\mu = \frac{1}{T} [2\eta\sigma_{\mu\nu}\sigma^{\mu\nu} + \zeta(\nabla_\rho u^\rho)^2]$$

## *Conclusions*


- effective dissipation can arise in quantum field theories due to effective local loss of information
- equations of motion for close-to-equilibrium theories can be obtained from analytic continuation
- general covariance and energy-momentum conservation lead to equations for fluid velocity and entropy production
- local form of second law of thermodynamics is implemented on the level of the effective action  $\Gamma[\Phi]$
- interesting applications

## Outlook

- proper understanding of local dissipation in terms of *entanglement entropy*  
[J. Berges, S. Floerchinger, R. Venugopalan, PLB 778 (2018) 442; JHEP 1804 (2018)145]



- causal dissipative relativistic fluid dynamics needs *hyperbolic* equations  
[S. Floerchinger, E. Grossi, 1711.06687]






*Backup slides*

## *Double time path formalism*

- formalism for general, far-from-equilibrium situations: Schwinger-Keldysh double time path
- can be formulated with two fields  $\Phi = \frac{1}{2}(\phi_+ + \phi_-)$ ,  $\chi = \phi_+ - \phi_-$
- in principle for arbitrary initial density matrices, in praxis mainly Gaussian initial states
- allows to treat also dissipation
- useful also to treat initial state fluctuations or forced noise in classical statistical theories
- difficult to recover thermal equilibrium, in particular non-perturbatively



## *Equations of motion from the Feynman action ?*

- consider damped harmonic oscillator as example. Time-ordered or Feynman action is obtained from analytic action by replacing  $s_1(\omega) \rightarrow \text{sign}(\omega)$

$$\Gamma_{\text{time ordered}}[x] = \int \frac{d\omega}{2\pi} \frac{m}{2} x^*(\omega) [-\omega^2 - 2i|\omega| \zeta\omega_0 + \omega_0^2] x(\omega)$$

- field equation  $\frac{\delta}{\delta x(t)} \Gamma_{\text{time ordered}}[x] = J(t)$  would give

$$[-\omega^2 - 2i|\omega| \zeta\omega_0 + \omega_0^2] x(\omega) = J(\omega)$$

- violates reality constraint  $x^*(\omega) = x(-\omega)$  for  $J^*(\omega) = J(-\omega)$
- solution not causal

$$x(t) = \int_{t'} \Delta_F(t - t') J(t')$$

because Feynman propagator  $\Delta_F(t - t')$  not causal.

- in contrast, retarded variation of analytic action leads to real and causal equation of motion

## Tree-like structures

- discontinuous terms in analytic action could be of the form

$$\Gamma_{\text{Disc}}[\Phi] = \int d^d x \sqrt{g} \left\{ f[\Phi](x) s_R(u^\mu(x) \frac{\partial}{\partial x^\mu}) g[\Phi](x) \right\}$$

- more general, tree-like structure are possible such as

$$\Gamma_{\text{Disc}}[\Phi] = \int_{x,y} \left\{ f[\Phi](x) s_R(u^\mu(x) \frac{\partial}{\partial x^\mu}) g[\Phi](x,y) s_R(u^\mu(y) \frac{\partial}{\partial y^\mu}) h[\Phi](y) \right\}$$

or

$$\Gamma_{\text{Disc}}[\Phi] = \int_{x,y,z} \left\{ f[\Phi](x) s_R(u^\mu(x) \frac{\partial}{\partial x^\mu}) g[\Phi](x,y,z) s_R(u^\mu(y) \frac{\partial}{\partial y^\mu}) h[\Phi](y) \right. \\ \left. \times s_R(u^\mu(z) \frac{\partial}{\partial z^\mu}) j[\Phi](z) \right\}$$

- for retarded variation calculate  $\delta\Gamma$  and set  $s_R(u^\mu \partial_\mu) \rightarrow -1$  if derivative operator points towards node that is varied and  $s_R(u^\mu \partial_\mu) \rightarrow 1$  if derivative operator points in opposite direction

## Analytic continuation of renormalization group equations

[Floerchinger, JHEP 1205 (2012) 021]

- consider a point  $p_0^2 - \vec{p}^2 = m^2$  where  $P_1(m^2) = 0$ .
- one can expand around this point


$$P_1 = Z(-p_0^2 + \vec{p}^2 + m^2) + \dots$$

$$P_2 = Z\gamma^2 + \dots$$

- leads to Breit-Wigner form of propagator (with  $\gamma^2 = m\Gamma$ )

$$G(p) = \frac{1}{Z} \frac{-p_0^2 + \vec{p}^2 + m^2 + i s(p_0) m\Gamma}{(-p_0^2 + \vec{p}^2 + m^2)^2 + m^2\Gamma^2}.$$

- a few parameters describe the singular structure of the propagator



## Truncation for relativistic scalar $O(N)$ theory

$$\Gamma_k = \int_{t, \vec{x}} \left\{ \sum_{j=1}^N \frac{1}{2} \bar{\phi}_j \bar{P}_\phi(i\partial_t, -i\vec{\nabla}) \bar{\phi}_j + \frac{1}{4} \bar{\rho} \bar{P}_\rho(i\partial_t, -i\vec{\nabla}) \bar{\rho} + \bar{U}_k(\bar{\rho}) \right\}$$

with  $\bar{\rho} = \frac{1}{2} \sum_{j=1}^N \bar{\phi}_j^2$ .

- Goldstone propagator massless, expanded around  $p_0 - \vec{p}^2 = 0$

$$\bar{P}_\phi(p_0, \vec{p}) \approx \bar{Z}_\phi (-p_0^2 + \vec{p}^2)$$

- radial mode is massive, expanded around  $p_0^2 - \vec{p}^2 = m_1^2$

$$\begin{aligned} \bar{P}_\phi(p_0, \vec{p}) + \bar{\rho}_0 \bar{P}_\rho(p_0, \vec{p}) + \bar{U}'_k + 2\bar{\rho} \bar{U}''_k \\ \approx \bar{Z}_\phi Z_1 \left[ (-p_0^2 + \vec{p}^2 + m_1^2) - is(p_0) \gamma_1^2 \right] \end{aligned}$$

## Flow of the effective potential

$$\partial_t U_k(\rho) \Big|_{\bar{p}} = \frac{1}{2} \int_{p_0=i\omega_n, \bar{p}} \left\{ \frac{(N-1)}{\bar{p}^2 - p_0^2 + U' + \frac{1}{Z_\phi} R_k} + \frac{1}{Z_1 [(\bar{p}^2 - p_0^2) - i s(p_0) \gamma_1^2] + U' + 2\rho U'' + \frac{1}{Z_\phi} R_k} \right\} \frac{1}{Z_\phi} \partial_t R_k.$$

- summation over Matsubara frequencies  $p_0 = i2\pi Tn$  can be done using contour integrals.
- radial mode has non-zero decay width since it can decay into Goldstone excitations.
- use Taylor expansion for numerical calculations

$$U_k(\rho) = U_k(\rho_{0,k}) + m_k^2(\rho - \rho_{0,k}) + \frac{1}{2} \lambda_k(\rho - \rho_{0,k})^2$$