
Charm and bottom flavored hadrons production from strangeness rich quark gluon plasma hadronization

Inga Kuznetsova and Johann Rafelski

Department of Physics, University of Arizona

We study QGP hadronization at given b, c quark content. We predict the yields of charm and bottom flavored hadrons within statistical hadronization model. The important new feature is that we take into account **high** strangeness and entropy content of QGP, conserving strangeness yield and entropy at hadronization.

The European Physical Journal C - Particles and Fields. C51, 113-133, (2007)
arXiv:hep-ph/0607203

Motivations

- Probe of QGP properties, confirmation of deconfinement
 - Information on hadronization temperature of heavy flavored hadrons
 - Understanding of properties of phase transition between deconfinement phase and hadronic gas (HG) phase in strangeness rich QGP.
-

Statistical model

Assumed Boltzmann distribution for b, c, s, hadrons:

$$\frac{dN_i}{dy} = \gamma_i n_i^{eq} \frac{dV}{dy}, \quad n_i^{eq} = \lambda_i \frac{T^3}{2\pi^2} g_i W(m_i / T),$$


$$\text{where } W(x) = x^2 K_2(x)$$

- $\lambda=1$ ($\mu = T \ln \lambda = 0$) for all particles
- γ_i is phase space occupancy factor

$\gamma_i = 1$, $n_i = n_i^{eq}$ is chemical equilibrium for particle i γ_i^Q is in QGP $i=c, b, s, q$ (q is u or d)

γ_i^H after hadronization, for example for D

$$\text{mesons: } \gamma_D^H = \gamma_c^H \gamma_q^H$$



Main model assumptions

- We do not assume chemical equilibrium for quark flavors.
- We work in framework of fast hadronization to final state. Physical conditions (system volume, temperature) do not change.

- Flavor conservation: $\frac{dN_i^{HG}}{dy} = \frac{dN_i^{QGP}}{dy}$

fixes statistical parameters ($\gamma_b^H, \gamma_c^H, \gamma_s^H$) for quark yield.

For LHC: $\frac{dN_c^{QGP}}{dy} = 10; \quad \frac{dN_b^{QGP}}{dy} = 1.$


- Entropy conservation: $\frac{dS^{QGP}}{dy} = \frac{dS^{HG}}{dy}$ fixes $\gamma_q^{HG} (\neq 1)$

- In QGP $\gamma_q^{QGP} = 1$

(The entropy of expanding QGP is conserved: $T^3 \frac{dV}{dy} \approx const.$)

Strangeness

- Strangeness (s) production in thermal gluon fusion follows in time entropy (S) production
- Ratio s/S depends on energy of collision, s increases faster with energy than S. The hot state, where the threshold for s production is exceeded, lives longer
- At RHIC energies $s/S \sim 0.03$, at LHC expect $0.03 \leq s/S \leq 0.05$



- obtained from SHARE 2.1

"SHARE: Statistical hadronization with resonances,"


G. Torrieri, S. Steinke, W. Broniowski, W. Florkowski, J. Letessier and J. R. Comput. Phys. Commun. 167, 229 (2005) (SHARE 1) [arXiv:nucl-th/0404083];

G. Torrieri, S. Jeon, J. Letessier and J. R. Comput. Phys. Commun. 175, 635 (2006) (SHARE 2) [arXiv:nucl-th/0603026]

Webpage:

<http://www.physics.arizona.edu/~torrieri/SHARE/sharev1.html>

Effect of strangeness on ratio D/D_s .





$$\frac{D}{D_s} = \left(\frac{\gamma_s^H}{\gamma_q^H} \right)^{-1} f(T)$$

$$D_s = \gamma_c \gamma_s n_{D_s}^{eq} = \gamma_c \gamma_s \sum_i n_{D_{s_i}}^{eq}$$


$$D = \gamma_c \gamma_q n_D^{eq} = \gamma_c \gamma_q \sum_i n_{D_i}^{eq}$$

	hadron		M[GeV]	hadron		M[GeV]	g
D	$D^0(0^-)$	$c\bar{u}$	1.8646	$B^0(0^-)$	$b\bar{u}$	5.279	1
	$D^+(0^-)$	$c\bar{d}$	1.8694	$B^+(0^-)$	$b\bar{d}$	5.279	1
	$D^{*0}(1^-)$	$c\bar{u}$	2.0067	$B^{*0}(1^-)$	$b\bar{u}$	5.325	3
	$D^{*+}(1^-)$	$c\bar{d}$	2.0100	$B^{*+}(1^-)$	$b\bar{d}$	5.325	3
	$D^0(0^+)$	$c\bar{u}$	2.352	$B^0(0^+)$	$b\bar{u}$	5.697	1
	$D^+(0^+)$	$c\bar{d}$	2.403	$B^+(0^+)$	$b\bar{d}$	5.697	1
	$D_1^{*0}(1^+)$	$c\bar{u}$	2.4222	$B_1^{*0}(1^+)$	$b\bar{u}$	5.720	3
	$D_1^{*+}(1^+)$	$c\bar{d}$	2.4222	$B_1^{*+}(1^+)$	$b\bar{d}$	5.720	3
	$D_2^{*0}(2^+)$	$c\bar{u}$	2.4589	$B_2^{*0}(2^-)$	$b\bar{u}$	(5.730)	5
	$D_2^{*+}(2^+)$	$c\bar{d}$	2.4590	$B_2^{*+}(2^+)$	$b\bar{d}$	(5.730)	5
D_s	$D_s^+(0^-)$	$c\bar{s}$	1.9868	$B_s^0(0^-)$	$s\bar{b}$	5.3696	1
	$D_s^{*+}(1^-)$	$c\bar{s}$	2.112	$B_s^{*0}(1^-)$	$s\bar{b}$	5.416	3
	$D_{sJ}^{*+}(0^+)$	$c\bar{s}$	2.317	$B_{sJ}^{*0}(0^+)$	$s\bar{b}$	(5.716)	1
	$D_{sJ}^{*+}(1^+)$	$c\bar{s}$	2.4593	$B_{sJ}^{*0}(1^+)$	$s\bar{b}$	(5.760)	3
	$D_{sJ}^{*+}(2^+)$	$c\bar{s}$	2.573	$B_{sJ}^{*0}(2^+)$	$s\bar{b}$	(5.850)	5

Ratio $D(B)/D_s(B_s)$ as a probe of T at measured s/S




Non-strange to strange charm baryons yields ratios as a function of γ_s/γ_q ratio



$$cqq / cqs \propto (\gamma_s / \gamma_q)^{-1}$$

$$cqq / css \propto (\gamma_s / \gamma_q)^{-2}$$

Double strange charm baryons (Ω_c^0) yield as a function of hadronization temperature T




Ω_c^0 (2700 MeV) decay modes:


- $\Sigma^+ K^- \pi^+$
- $\Xi^0 K^- \pi^+$
- $\Xi^- K^- \pi^+ \pi^+$
- $\Omega^- \pi^+$
- $\Omega^- \pi^+ \pi^0$
- $\Omega^- \pi^- \pi^+ \pi^+$

Total yield of all hidden charm mesons as a function of T.

hadron		mass(GeV)	g
$\eta_c(1S)$	$c\bar{c}$	2.9779	1
$J/\Psi(1S)$	$c\bar{c}$	3.0970	3
$\chi_{c0}(1P)$	$c\bar{c}$	3.4152	1
$\chi_{c1}(1P)$	$c\bar{c}$	3.5106	3
$h_c(1P)$	$c\bar{c}$	3.526	3
$\chi_{c2}(1P)$	$c\bar{c}$	3.5563	5
$\eta_c(2S)$	$c\bar{c}$	3.638	1
$\psi(2S)$	$c\bar{c}$	3.686	3
ψ	$c\bar{c}$	3.770	3
ψ	$c\bar{c}$	3.836	5
ψ	$c\bar{c}$	4.040	3
ψ	$c\bar{c}$	4.159	3
ψ	$c\bar{c}$	4.415	3



J/ Ψ yield as a function of γ_s/γ_q




- Both entropy and strangeness contents enhancement may result to J/ Ψ suppression.
- More light and/or strange quarks more probability for charm to bound to these quarks than to find anti-charm quark.

Conclusions

- Phase space occupancy factors of strange and light quarks have strong influence on heavy flavor hadron production.
- Significant increase of the yield of strange quark-containing charm (bottom) mesons and baryons with increase of s/S as compared to the chemical equilibrium yields.
- The change in the yield of hadrons without strangeness but with light quark(s) depends on both s/S and γ_q . The ratio of these hadrons to similar strange hadrons always decreases with increase of s/S .
- Yields of hadrons with two heavy quarks, as J/Ψ , decrease compared to chemical equilibrium when γ_q and/or $\gamma_s > 1$. This may provide a mechanism of J/Ψ suppression.

Entropy after hadronization


- Because of liberation of color degree of freedom
$$\sigma^Q \geq 3\sigma^H$$
- The excess of entropy is observed in the multiplicity of particles in final state.
- After hadronization $S^Q \approx S^H$, $\gamma_q^H > 1$.
- When $\gamma_q^H = \gamma_q^{cr}$: Bose singularity for pions; Maximum of possible entropy content after hadronization




Strangeness conservation during hadronization

$$s = \frac{dV}{dy} \left[\gamma_s^H \left(\gamma_q^H n_K^{eq} + \gamma_q^{H2} n_Y^{eq} \right) + 2\gamma_s^{H2} \gamma_q^H n_{\Xi}^{eq} \right]$$

- The equilibrium densities n_i^{eq} are sums of all known states densities for given particle i .



Charm (bottom) hadronization



- c,b quarks produced in first nn collisions.
- $c=10$, $b=1$;
- Flavor conservation equation


$$c = \frac{dV}{dy} \left[\gamma_c^H n_{open}^c + \gamma_c^{H2} \left(n_{hid}^c + 2\gamma_q^H n_{ccq}^{eq} + 2\gamma_s^H n_{ccs}^{eq} \right) \right];$$

$$n_{open}^c = \gamma_q^H n_D^{eq} + \gamma_s^H n_{Ds}^{eq} + \gamma_q^{H2} n_{qqc}^{eq} + \gamma_s^H \gamma_q^H n_{sqc}^{eq} + \gamma_q^{H2} n_{ssc}^{eq};$$

$$n_{hid}^c = \gamma_c^{H2} n_{\bar{c}c}^{eq}.$$

- $\gamma_b^H \gg \gamma_c^H \gg \gamma_s^H$
- Equilibrium case when $\gamma_q^H = \gamma_s^H = 1$

D(B), D_s(B_s) mesons yield as a function of γ_s/γ_q .



$$D(B) / N_{c(b)} \propto \gamma_{c(b)} \gamma_q / N_{c(b)}$$

$$D_s(B_s) / N_{c(b)} \propto \gamma_{c(b)} \gamma_s / N_{c(b)}$$

$\gamma_{c(b)}/N_{c(b)}$ is almost independent from $N_{c(b)}$.