

Exclusive processes at COMPASS – present and future

Materials for discussion

Andrzej Sandacz

Sołtan Institute for Nuclear Studies, Warsaw


workshop on ‘Hard Exclusive Process at JLab 12 GeV and a Future EIC’



University of Maryland College Park, October 29-30, 2006

COMPASS experiment

~ 250 physicists
from 28 institutes




μ 160 GeV
from CERN SPS
Beam Polarization ~ -76%

Beam intensity
Luminosity

$2 \cdot 10^8 \mu^+/\text{spill}$ (4.8 s / 16.2 s)
 $\sim 5 \cdot 10^{32} \text{ cm}^{-2}\text{s}^{-1}$

Incoherent exclusive ρ^0 production



^6LiD polarized target


Kinematics:

$$\nu > 30 \text{ GeV}$$

$$E_{\mu'} > 20 \text{ GeV}$$

$$Q^2 > 0.01 \text{ GeV}^2$$

(Q^2 cut applied only in SDME analysis)



Assuming
both hadrons are π
 $0.5 < M_{\pi\pi} < 1 \text{ GeV}$

Exclusivity of
the reaction


$$E_{\text{miss}} = (M_X^2 - M_N^2) / 2M_N$$

$-2.5 < E_{\text{miss}} < 2.5 \text{ GeV}$

Incoherent production
 $0.15 < p_T^2 < 0.5 \text{ GeV}^2$
scattering off a
quasi-free nucleon

Background $\sim 12\%$

Kinematical domain of the final sample





2002 : 800,000 evts



2003 : 1,600,000 evts

+ 2004 : not yet analyzed
~ will double the data sample

$$\langle p_{\perp}^2 \rangle = 0.27 \text{ GeV}^2$$



COMPASS results on A_1^ρ (d)



COMPASS results on A_1^ρ on polarized deuteron target consistent with 0


Extended kinematical range of COMPASS by almost 2 decades down both in Q^2 and x

COMPASS : inner bars –stat. outer – total errors
HERMES: total errors

➔ at small x (< 0.01) contribution of unnatural parity exchanges small for exclusive ρ^0 production

Measurement of r_{00}^{04}

$0.01 < Q^2 < 0.05 < Q^2 < 0.3 < Q^2 < 0.6 < Q^2 < 2.0 < Q^2 < 10 \text{ GeV}^2$



Distribution :

$$W(\cos\theta) = \frac{3}{4} \left[(1 - r_{00}^{04}) + (3r_{00}^{04} - 1)\cos^2\theta \right]$$


Spin density matrix element:

$$r_{00}^{04} = \frac{|T_{01}|^2 + (\varepsilon + \delta)|T_{00}|^2}{N_T (1 + (\varepsilon + \delta)R)} \xrightarrow{\text{SCHC}} \frac{\sigma_L}{\sigma_T}$$

$$R = \sigma_L / \sigma_T \quad N_T = |T_{11}|^2 + |T_{-11}|^2 + |T_{01}|^2$$

$$\Gamma_L / \Gamma_T = \varepsilon + \delta$$

$T_{\lambda\mu\lambda\gamma}$ helicity amplitudes
 meson photon



Determination of $R = \sigma_L/\sigma_T$


If SCHC holds :

only $T_{00} \neq 0$
 $T_{11} \neq 0$

Then :


$$R = \frac{\sigma_L}{\sigma_T} = \frac{1}{(\varepsilon + \delta)} \frac{r_{00}^{04}}{1 - r_{00}^{04}}$$

Impact on GPD study:
determination of σ_L
 σ_L is dominant at $Q^2 > 2 \text{ GeV}^2$



- High statistics from quasi-photoproduction to hard production
- Better coverage at high Q^2 with 2003 and 2004 data

Measurement of r_{1-1}^{04} and $\text{Im } r_{1-1}^3$



Distribution :



$$W(\varphi) = \frac{1}{2\pi} [1 - 2r_{1-1}^{04} \cos 2\varphi + 2 \text{Im} r_{1-1}^3 P_\mu \sqrt{1 - \varepsilon^2} \sin 2\varphi]$$

beam polarisation

Spin density matrix elements:

$$r_{1-1}^{04} = \frac{\text{Re}(T_{11} T_{-11}^*) - (\varepsilon + \delta) |T_{10}|^2}{N_T (1 + (\varepsilon + \delta) R)} = 0$$


$$\text{Im} r_{1-1}^3 = \dots = 0 \quad \leftarrow \text{If SCHC holds}$$



Summary and outlook for vector mesons

- Asymmetry $A_1^p(d)$ consistent with zero over wide range of Q^2 and x
small contribution of unnatural parity exchanges at $x < 0.01$
first measurement at small Q^2 and small x
- High-statistics data on **SDM elements and R** for incoherent exclusive ρ^0 production
in a wide Q^2 range (including small Q^2 not covered previously)
weak violation of SCHC observed
- Significant improvement of accuracy expected after including 2004 (and 2003 data)
extraction of **23 SDMEs** under way
- Studies of **coherent** exclusive ρ^0 production foreseen
- Single spin asymmetry for **transversely polarized target** \rightarrow **E/H** GPDs
- Analysis of exclusive ϕ and J/ψ production in progress

possible solution to complete the COMPASS setup




2004-2007:


Received funding by EU FP6 (Bonn-Mainz-Warsaw-Saclay)

Goal: full test of feasibility of a 45° sector recoil detector

- scintillating material studies (200ps ToF Resolution over 4m)
- fast triggering and multi-hit ADC/TDC system



DVCS+ Bethe Heitler




BH calculable

The high energy muon beam at COMPASS allows to play with the relative contributions DVCS-BH which depend on



$$1/\gamma = 2 m_p E_\ell x_{Bj} / Q^2$$

Higher energy: DVCS \gg BH
 \Rightarrow DVCS Cross section

Smaller energy: DVCS \sim BH
 \Rightarrow Interference term will provide the DVCS amplitude



$Q^2 = 4 \text{ GeV}^2$



Advantage of $\vec{\mu}^+$ and $\vec{\mu}^-$ for Deeply virtual Compton scattering (+Bethe-Heitler)

$$A_{(\mu p \rightarrow \mu p \gamma)}^{DVCS} = \int_{-1}^{+1} dx \frac{H(x, \xi, t)}{x - \xi + i\epsilon} = \mathcal{P} \int_{-1}^{+1} dx \frac{H(x, \xi, t)}{x - \xi} - i\pi H(x = \xi, \xi, t)$$

$t, \xi \sim x_{Bj/2}$ fixed

$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = \cancel{d\sigma^{BH} + d\sigma^{DVCS}_{unpol}} + \cancel{P_{\mu} d\sigma^{DVCS}_{pol}}$$


$$+ \cancel{e_{\mu} a^{BH} \text{Re} A^{DVCS}}$$

$$\times \cos n\varphi$$

$$+ \cancel{P_{\mu} d\sigma^{DVCS}_{pol}}$$

$$+ \cancel{e_{\mu} P_{\mu} a^{BH} \text{Im} A^{DVCS}}$$

$$\times \sin n\varphi$$



$$P_{\mu^+} = -0.8 \quad P_{\mu^-} = +0.8$$

$$\sigma^{\vec{\mu}^+} + \sigma^{\vec{\mu}^-} \sim H(x = \xi, \xi, t)$$


$$\sigma^{\vec{\mu}^+} - \sigma^{\vec{\mu}^-} \sim \mathcal{P} \int_{-1}^{+1} dx \frac{H(x, \xi, t)}{x - \xi}$$

COMPASS

6 angular distributions


among 18: 3 bins in $x_{Bj}=0.05, 0.1, 0.2$
6 bins in Q^2 from 2 to 7 GeV^2

BCA $E_\mu=100GeV$ $\vartheta=1deg$



BCA in DVCS
projections
for 1 year

HERMES



if $N_\mu \times 5 \Rightarrow Q^2 < 17 \text{ GeV}^2$
for DVCS

Benefit of a higher
muon intensity
for GPDs study

if $N_\mu \times 2 \Rightarrow Q^2 < 11 \text{ GeV}^2$
for DVCS

Limitation by luminosity

now $N_\mu = 2 \cdot 10^8 \mu$ per SPS spill
for DVCS
 $\Rightarrow Q^2 < 7.5 \text{ GeV}^2$

At fixed x_{Bj} , study in Q^2

