

Low Energy Neutrino Physics

Nikolai Tolich

Outline

- Double-beta-decay
- Geoneutrinos
- Solar neutrinos

Beta-decay


- β^- -decay is the conversion of a bound neutron into a proton with the emission of an electron and an electron anti-neutrino




- β^+ -decay is the conversion of a bound proton into a neutron with the emission of a positron and an electron neutrino



- Electron-capture is similar to β^+ -decay but instead of producing a positron, an electron is captured




Conservation of energy




- Conservation of energy tells us that for
 - β -decay:
mass of the X is greater than the sum of the mass of Y, the electron, and neutrino
 - Electron-capture:
mass of the X plus electron is greater than the mass of Y plus the neutrino
- Nuclear physicists often draw energy level diagrams for a particular atomic mass, like that on the left

Double-beta-decay




- Double-beta-decay occurs when the single beta-decay is energetically forbidden.
- It is an extremely rare process, but has been observed in numerous isotopes.



Neutrinoless double-beta-decay


- If neutrinos are Majorana particles (an anti-neutrino is equivalent to a neutrino) then we do not need to emit any neutrinos in a double-beta-decay.
- Neutrinoless double-beta-decay is the best probe we have to test if neutrinos are Majorana particles.



Majorana neutrinos?

- So-called “seesaw” models, which explain the lightness of the neutrino, require the neutrino to be a Majorana particle.
- Leptogenesis, which could explain the apparent matter dominance of the universe also requires neutrinos to be a Majorana particle.
- **All other particles are Dirac particles.**


Observing neutrinoless double-beta-decay




- Because there are no neutrinos to take away energy, the total electron energy in neutrinoless double-beta-decay is equal to the decay energy.
- We can recognize neutrinoless double-beta-decay by observing a peak at the end of the total electron energy spectrum.

SNO (SNO+) detector

- Located in the Vale Inco Ltd. Creighton Mine near Sudbury, Canada
 - 1 kton D₂O held in 12 m diameter acrylic vessel, **to be replaced with Nd doped liquid scintillator**
 - 18 m diameter support structure holds 9500 PMTs (~60% photocathode coverage)
 - 1.7 kton inner shielding H₂O
 - 5.3 ktons outer shielding H₂O



Expected energy spectrum in SNO+



Geoneutrinos

Convection in Earth





Image: by Surachit, http://en.wikipedia.org/wiki/File:Oceanic_spreading.svg


- Seismic data splits Earth into 5 basic regions:
 - inner core
 - outer core
 - mantle
 - oceanic crust
 - continental crust
- All these regions are solid except the outer core.
- The mantle convects even though it is solid.
- Oceanic crust is being renewed at mid-ocean ridges and recycled at trenches.

Heat flow from the Earth

Bore-hole measurements



Heat flow




- Conductive heat flow measured from bore-hole temperature gradient and conductivity
- Total heat flow 46 ± 2 TW
- Based on chondritic meteorites the heat production from U, Th, and K are 8 TW, 8 TW, and 3 TW, respectively.

Image: Pollack et. al

Discrepancy?

- The measured total heat flow is **44 TW**.
- The estimated radiogenic heat produced is **19 TW**.
- Models of mantle convection suggest that the radiogenic heat production rate should be a large fraction of the measured heat flow.
- Problem with
 - Mantle convection model?
 - Total heat flow measured?
 - Estimated amount of radiogenic heat production rate?
- Geoneutrinos can serve as a cross-check of the radiogenic heat production.

Geo-neutrino signal



A little history



Fred Reines (circa 1953)

Dear Fred,
Just occurred to me
that your background
neutrinos may just be comming
from high energy β -decaying
members of U and Th families
in the crust of the Earth. We
not have on the train any
inform. to check it up, but it
seems the order of magn. is
reasonable. In fact the total energy
radioactive energy production
under one square foot of surface
may well be equal to the
energy of solar radiation falling
on ~~Earth~~ that surface...
what do you think?
Write to me at: The Union
Univ. of Mich. Ann Arbor. Mich
Yours G-CO.



Were they geo-neutrinos?

TO: DR. GEORGE GAMOW
THE UNION
UNIVERSITY OF MICHIGAN
ANN ARBOR, MICHIGAN

MESSAGE:

FROM NUMBERS IN VREY BOOK ON THE PLANETS, EQUILIBRIUM HEAT LOSS
FROM EARTH'S SURFACE IS $50 \text{ ERGS}/\text{CM}^2 \text{ SEC.}$ IF ASSUME ALL DUE TO
BETA DECAY THEN HAVE ONLY ENOUGH ENERGY FOR ABOUT 10^8 , 10^{11} Mev
NEUTRONS PER CM^2 AND SEC. THIS IS LOW BY 10^5 OR SO. SHORT
HALF LIVES WOULD BE MADE BY COSMIC RAYS OR NEUTRONS IN EARTH.
IN VIEW OF RARITY OF COSMIC RAYS: I.E. ABOUT EQUAL TO ENERGY
OF STARLIGHT AND OF NEUTRONS IN EARTH THIS SOURCE OF NEUTRONS
SEEMS EVEN LESS LIKELY AS A SOURCE OF OUR SIGNAL.

S/N crust and mantle




Image: S. Enomoto

S/N Mantle






Image: S. Enomoto

Where do the neutrinos come from?



Detecting electron anti neutrinos

- $\bar{\nu}_e + p \rightarrow e^+ + n$
The positron energy is related to the neutrino energy.
- The positron loses its energy then annihilates with an electron.
- The neutron first thermalizes then is captured by a proton with a mean capture time of $\sim 200\mu s$.




Need for more detectors


- Current geoneutrino flux measurements
 - KamLAND $(4.3^{+1.2}_{-1.1}) \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$
 - BOREXINO $(7.1^{+2.9}_{-2.4}) \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$
- Ideally we would like to make a measurement in the ocean to probe the contribution from the mantle.
- However, with a precision measurement on the crust, and an accurate measurement of the local crustal contribution from heat flow measurements, could also obtain mantle contribution.
- Ideally we would like multiple measurements to probe local variations (despite our assumptions, the mantle is not uniform).

Solar neutrinos

Solar pp chain reactions



Neutrino energy spectrum




Ray Davis




- $^{37}\text{Cl} + \nu_e \rightarrow ^{37}\text{Ar} + e^-$
- ^{37}Ar is a gas which is removed from detector with He carrier gas
- Outside the active volume the ^{37}Ar is detected via $^{37}\text{Al} + e^- \rightarrow ^{37}\text{Cl} + \nu_e$ which has a half-life of 35 days

SuperK








- 50,000 ton ring-imaging water Cherenkov detector
- SuperK detects solar neutrinos from electron elastic scattering



- $\sigma(\nu_e) \approx 6 \sigma(\nu_\mu) \approx 6 \sigma(\nu_\tau)$
- Strong directionality
- The scattered electron produces a Cherenkov ring


SNO: 3 reactions and 3 phases

- Detected neutrons produced in NC reactions three ways
- Phase I:
 - Signal observed in PMT array via $n + {}^2H \rightarrow {}^3H + 6.25\text{ MeV}$
- Phase II:
 - Added salt
 - Signal observed in PMT array via $n + {}^{35}Cl \rightarrow {}^{36}Cl + 8.6\text{ MeV}$
- Phase III:
 - Removed salt and added an array of 40 3He filled proportional counters “neutral-current detectors” (NCDs).
 - Signal observed in NCD array via $n + {}^3He \rightarrow {}^3H + p$




Solar neutrino measurements


Total Rates: Standard Model vs. Experiment
Bahcall–Serenelli 2005 [BS05(OP)]




Theory ■ ^7Be ■ p-p, pep ■ Experiments ■
 ■ ^8B ■ CNO Uncertainties ■




Pretending to be miners



Very clean miners



Super-K cleaning



Boating in Super-K, not likely again



Questions?