

# THE UNITARY GAS: SYMMETRY PROPERTIES AND APPLICATIONS

Yvan Castin, Félix Werner, Christophe Mora

LKB and LPA, Ecole normale supérieure (Paris, France)

Ludovic Pricoupko

LPTMC, Université Paris 6



## OUTLINE OF THE TALK

- What is the unitary gas ?
- Simple facts from scaling invariance
- Time-dependent solution in a trap
- Separability in hyperspherical coordinates
- The 4-body Efimov effect

**WHAT IS THE UNITARY GAS ?**

## DEFINITION OF THE UNITARY GAS

- Non-relativistic particles with  $s$ -wave binary interaction with a two-body scattering amplitude

$$f_k = -\frac{1}{ik} \quad \forall k$$

- “Maximally” interacting: Unitarity of  $S$  matrix imposes  $|f_k| \leq 1/k$ .
- In real experiments with magnetic Feshbach resonance (Thomas, Salomon, Jin, Ketterle, Grimm, ...) :

$$-\frac{1}{f_k} = \frac{1}{a} + ik - \frac{1}{2}k^2 r_e + O(k^4 b^3)$$

almost unitary if “infinite” scattering length  $a$  and “zero” ranges:

$$k_{\text{typ}}|a| \gg 1, k_{\text{typ}}|r_e| \ll 1, k_{\text{typ}}b \ll 1.$$

- All these two-body conditions are only necessary.

# THE ZERO-RANGE WIGNER-BETHE-PEIERLS MODEL

- Interactions are replaced by contact conditions.
- For  $r_{ij} \rightarrow 0$  with fixed  $ij$ -centroid  $\vec{C}_{ij} = (m_i\vec{r}_i + m_j\vec{r}_j)/(m_i + m_j)$  different from  $\vec{r}_k, k \neq i, j$ :

$$\psi(\vec{r}_1, \dots, \vec{r}_N) = \left( \frac{1}{r_{ij}} - \frac{1}{a} \right) A_{ij}[\vec{C}_{ij}; (\vec{r}_k)_{k \neq i, j}] + O(r_{ij})$$

- Elsewhere, non interacting Schrödinger equation

$$E\psi(\vec{X}) = \left[ -\frac{\hbar^2}{2m} \Delta_{\vec{X}} + \frac{1}{2}m\omega^2 X^2 \right] \psi(\vec{X})$$

with  $\vec{X} = (\vec{r}_1, \dots, \vec{r}_N)$ .

- Exchange symmetry: Even for boson positions, odd for same-spin fermion positions.
- Unitary gas exists iff Hamiltonian is self-adjoint.

# **SIMPLE FACTS FROM SCALING INVARIANCE**

# SCALING INVARIANCE OF CONTACT CONDITIONS

$$\psi(\vec{X}) \underset{r_{ij} \rightarrow 0}{=} \frac{1}{r_{ij}} A_{ij}[\vec{C}_{ij}; (\vec{r}_k)_{k \neq i,j}] + O(r_{ij})$$

- Domain of Hamiltonian is scaling invariant: If  $\psi$  obeys the contact conditions, so does  $\psi_\lambda$  with

$$\psi_\lambda(\vec{X}) \equiv \frac{1}{\lambda^{3N/2}} \psi(\vec{X}/\lambda)$$

- Consequences (also true for the ideal gas):


free space	box (periodic b.c.)	trap
$\forall N$ , no bound states <sup>(*)</sup>	$PV = 2E/3$	virial $E = 2E_{\text{harm}}$

<sup>(\*)</sup> If  $\psi$  of eigenenergy  $E$ ,  $\psi_\lambda$  of eigenenergy  $E/\lambda^2$ . Square integrable eigenfunctions (after center of mass removal) correspond to point-like spectrum, for selfadjoint  $H$ .

## USEFUL CONSTRAINTS FOR MONTE CARLO

$$[\xi = \mu(T = 0)/E_F \leq 0, 41 \text{ (Carlson, 2009)}]$$

Burovski, Prokof'ev, Svistunov, Troyer (2006),  $T_c/T_F = 0.152(7)$ ; Goulko, Wingate (2010),  $T_c/T_F = 0.173(6)$





# TIME-DEPENDENT SOLUTION IN A TRAP

## IN A TIME-DEPENDENT TRAP

- At  $t = 0$  : static trap  $U(\mathbf{r}) = m\omega^2 r^2/2$ , system in eigenstate  $\psi_0(\vec{X})$  of energy  $E$ .
- For  $t > 0$ , arbitrary time dependence of trap spring constant,  $\omega(t)$ . Known solution for ideal gas:

$$\psi(\vec{X}, t) = \frac{e^{-i\theta(t)}}{\lambda^{3N/2}(t)} \exp\left[\frac{im\dot{\lambda}}{2\hbar\lambda} X^2\right] \psi_0(\vec{X}/\lambda(t))$$

with  $\ddot{\lambda} = \omega^2\lambda^{-3} - \omega^2(t)\lambda$  and  $\dot{\theta} = E\lambda^{-2}/\hbar$ .

- This is a gauge plus scaling transform.
- The gauge transform also preserves contact conditions:

$$r_i^2 + r_j^2 = 2C_{ij}^2 + \frac{1}{2}r_{ij}^2$$

so solution also applies to unitary gas!

Y. Castin, *Comptes Rendus Physique* 5, 407 (2004).

## IN THE MACROSCOPIC LIMIT

$$\psi(\vec{X}, t) = \frac{e^{-i\theta(t)}}{\lambda^{3N/2}} \exp \left[ \frac{im\dot{\lambda}}{2\hbar\lambda} X^2 \right] \psi_0(\vec{X}/\lambda)$$

density $\rho(\vec{r}, t) = \rho_0(\vec{r}/\lambda)/\lambda^3$	velocity field $\vec{v}(\vec{r}, t) = \vec{r} \dot{\lambda}/\lambda$
local temp. $T(\vec{r}, t) = T/\lambda^2$	pressure $P(\vec{r}, t) = P_0(\vec{r}/\lambda)/\lambda^5$
local entropy per particle	$s(\vec{r}, t) = s_0(\vec{r}/\lambda)$

This has to solve the hydrodynamic equations for a normal gas. Entropy production equation:

$$\rho k_B T (\partial_t s + \vec{v} \cdot \vec{\nabla} s) = \vec{\nabla} \cdot (\kappa \nabla T) + \zeta (\vec{\nabla} \cdot \vec{v})^2 + \frac{\eta}{2} \sum_{i,j} \left( \frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} - \frac{2}{3} \delta_{ij} \vec{\nabla} \cdot \vec{v} \right)^2$$

so the bulk viscosity is zero:  $\zeta(\rho, T) = 0 \forall T > T_c$ . Reproduces the conformal invariance result of Son (2007).

## LADDER STRUCTURE OF THE SPECTRUM

- Infinitesimal change of  $\omega$  for  $0 < t < t_f$ . For  $t > t_f$ :

$$\lambda(t) - 1 = \epsilon e^{-2i\omega t} + \epsilon^* e^{2i\omega t} + O(\epsilon^2)$$

so an undamped mode of frequency  $2\omega$ .

- Corresponding wavefunction change:

$$\psi(\vec{X}, t) = \left[ e^{-iEt/\hbar} - \epsilon e^{-i(E+2\hbar\omega)t/\hbar} L_+ + \epsilon^* e^{-i(E-2\hbar\omega)t/\hbar} L_- \right] \psi_0(\vec{X}) + O(\epsilon^2)$$


- Raising and lowering operators:

$$L_{\pm} = \pm i \left[ \frac{3N}{2i} - i\vec{X} \cdot \partial_{\vec{X}} \right] + \frac{H}{\hbar\omega} - m\omega X^2/\hbar$$

(in red, generator of scaling transform)

- Spectrum=collection of semi-infinite ladders of step  $2\hbar\omega$ .  
 $SO(2, 1)$  hidden symmetry (Pitaevskii, Rosch, 1997).

# LADDER STRUCTURE OF THE SPECTRUM (2)



## A USEFUL MAPPING

- Each energy ladder has a ground step of energy  $E_g$ , eigenfunction  $\psi_g$ .
- Integration of  $L_- \psi_g = 0$  gives, with  $\vec{X} = X \vec{n}$ :

$$\psi_g(\vec{X}) = e^{-m\omega X^2/2\hbar} X^{E_g/(\hbar\omega) - 3N/2} f(\vec{n})$$

- Limit  $\omega \rightarrow 0$  : mapping to zero energy free space solutions. N.B.:  $E_g/(\hbar\omega)$  is a constant.
- Free space problem solved for  $N = 3$  (Efimov, 1972)... so trapped case also solved (Werner, Castin, 2006).

# SEPARABILITY IN HYPERSPHERICAL COORDINATES

# SEPARABILITY IN HYPERSPHERICAL COORDINATES

Werner, Castin (2006)

- Use Jacobi coordinates to separate center of mass  $\vec{C}$
- Hyperspherical coordinates (arbitrary masses  $m_i$ ):

$$(\vec{r}_1, \dots, \vec{r}_N) \leftrightarrow (\vec{C}, R, \vec{\Omega})$$

with  $3N - 4$  hyperangles  $\vec{\Omega}$  and the hyperradius

$$m_u R^2 = \sum_{i=1}^N m_i (\vec{r}_i - \vec{C})^2$$

where  $m_u$  is an arbitrary mass unit.

- Hamiltonian is clearly separable:

$$H_{\text{internal}} = -\frac{\hbar^2}{2m_u} \left[ \partial_R^2 + \frac{3N-4}{R} \partial_R + \frac{1}{R^2} \Delta_{\vec{\Omega}} \right] + \frac{1}{2} m_u \omega^2 R^2$$



## Do the contact conditions preserve separability ?

- For free space  $E = 0$ , yes, due to scaling invariance:

$$\psi_{E=0} = R^{s_N - (3N-5)/2} \phi(\vec{\Omega}).$$

$E = 0$  Schrödinger's equation implies

$$\Delta_{\vec{\Omega}} \phi(\vec{\Omega}) = - \left[ s_N^2 - \left( \frac{3N-5}{2} \right)^2 \right] \phi(\vec{\Omega})$$

with contact conditions.  $s_N^2 \in$  discrete real set.

- For arbitrary  $E$ , Ansatz with  $E = 0$  hyperrangular part obeys contact conditions [ $R^2 = R^2(r_{ij} = 0) + O(r_{ij}^2)$ ]:

$$\psi = F(R) R^{-(3N-5)/2} \phi(\vec{\Omega})$$

- Schrödinger's equation for a fictitious particle in 2D:

$$EF(R) = -\frac{\hbar^2}{2m_u} \Delta_R^{2D} F(R) + \left[ \frac{\hbar^2 s_N^2}{2m_u R^2} + \frac{1}{2} m_u \omega^2 R^2 \right] F(R)$$

## SOLUTION OF HYPERRADIAL EQUATION ( $N \geq 3$ )

$$EF(R) = -\frac{\hbar^2}{2m_u} \Delta_R^{2D} F(R) + \left[ \frac{\hbar^2 s^2}{2m_u R^2} + \frac{1}{2} m_u \omega^2 R^2 \right] F(R)$$

- Which boundary condition for  $F(R)$  in  $R = 0$ ? Wigner-Bethe-Peierls does not say.
- Key point: particular solutions  $\sim R^{\pm s}$  for  $R \rightarrow 0$ .

$s > 1$	$0 < s < 1$	$s \in i\mathbb{R}^{+*}$
$F \sim R^s$	$F \sim (qR)^s \pm (qR)^{-s}$	$F \sim \text{Im} [(qR)^s]$
0 bound st.	one bound st. if –	$\infty$ nber of bound st.
$E_n = (2n + s + 1)\hbar\omega, n \geq 0$	$E \propto -\frac{\hbar^2 q^2}{m_u} :$ <b><math>N</math>-body resonance</b>	$E_n \propto -\frac{\hbar^2 q^2}{m_u} e^{-2\pi n/ s },$ $n \in \mathbb{Z} : \text{Efimov effect}$

# THE 4-BODY EFIMOV EFFECT

## THREE-BODY EFIMOV EFFECT

- Efimov (1971): Three bosons,  $1/a = 0$ , no dimer state. Then there exists an infinite number of trimer states,  $E = 0$  accumulation point, geometric spectrum:

$$E_n^{(3)} \underset{n \rightarrow +\infty}{\sim} E_{\text{ref}}^{(3)} e^{-2\pi n/|s_3|}$$

where purely imaginary  $s_3 = i \times 1.00624$  solves transcendental equation,  $E_{\text{ref}}^{(3)}$  depends on microscopic details.

- Efimov (1973): Solution for three arbitrary particles,  $1/a = 0$ . E.g. Efimov trimers for two fermions (masse  $M$ , same spin state) and one impurity (masse  $m$ ) if (Petrov, 2003)

$$\alpha \equiv \frac{M}{m} > \alpha_c(2; 1) \simeq 13.607$$

with  $s_3(\alpha) \in i\mathbb{R}^{+*}$  from known transcendental equation.

## ARE THERE EFIMOVIAN TETRAMERS ?

$$E_n^{(4)} \underset{n \rightarrow +\infty}{\sim} E_{\text{ref}}^{(4)} e^{-2\pi n/|s_4|} ?$$

### Negative results:

- Amado, Greenwood (1973): “There is No Efimov effect for Four or More Particles”. Explanation: Case of bosons, there exist trimers, tetramers decay.
- Hammer, Platter (2007), von Stecher, D’Incao, Greene (2009), Deltuva (2010): The four-boson problem (here  $1/a = 0$ ) depends only on  $E_{\text{ref}}^{(3)}$ , no  $E_{\text{ref}}^{(4)}$  to add.
- Key point:  $N = 3$  Efimov effect breaks separability in hyperspherical coordinates for  $N = 4$ .

Idea: Consider three fermions ( $M$ ) and one impurity ( $m$ ).

## REMINDER: MAIN POINTS OF GENERAL THEORY

- To find  $N$ -body Efimov effect, one simply needs to calculate the exponents  $s_N$ , that is to solve the Wigner-Bethe-Peierls model at zero energy:

$$\psi_{E=0}(\vec{r}_1, \dots, \vec{r}_N) = R^{s_N - (3N-5)/2} \phi(\vec{\Omega})$$

- The  $N$ -body Efimov effect takes place if and only if one of the  $s_N^2$  is  $< 0$ .
- General theory OK if  $\Delta_{\vec{\Omega}}$  self-adjoint: no  $n$ -body Efimov effect  $\forall n \leq N - 1$ .

## THE 3 + 1 FERMIONIC PROBLEM (Castin, Mora, Pricoupenko, 2010)

- Three fermions (mass  $M$ , same spin state) and one impurity (mass  $m$ )
- General theory OK for a mass ratio

$$\alpha \equiv \frac{M}{m} < \alpha_c(2; 1) \simeq 13.607$$

- Calculate  $E = 0$  solution in momentum space. An integral equation for Fourier transform of  $A_{ij}$ :

$$0 = \left[ \frac{1 + 2\alpha}{(1 + \alpha)^2} (k_1^2 + k_2^2) + \frac{2\alpha}{(1 + \alpha)^2} \vec{k}_1 \cdot \vec{k}_2 \right]^{1/2} D(\vec{k}_1, \vec{k}_2) + \int \frac{d^3 k_3}{2\pi^2} \frac{D(\vec{k}_1, \vec{k}_3) + D(\vec{k}_3, \vec{k}_2)}{k_1^2 + k_2^2 + k_3^2 + \frac{2\alpha}{1+\alpha} (\vec{k}_1 \cdot \vec{k}_2 + \vec{k}_1 \cdot \vec{k}_3 + \vec{k}_2 \cdot \vec{k}_3)}$$

- $D$  has to obey fermionic symmetry.

# REDUCTION OF THE INTEGRAL EQUATION

## Rotational invariance:

- $D$  is the  $m_l = 0$  component of a spinor of spin  $l$ :

$$\vec{D}(\vec{k}_1, \vec{k}_2) = {}^t \rho \vec{D}(\mathcal{R}\vec{k}_1, \mathcal{R}\vec{k}_2)$$

- Clever choice of the rotation matrix  $\mathcal{R}$ :

$$\vec{D}(\vec{k}_1, \vec{k}_2) = {}^t \rho \underbrace{\vec{D}[k_1 \vec{e}_x, k_2 (\cos \theta \vec{e}_x + \sin \theta \vec{e}_y)]}_{2l+1 \text{ unknown functions } f_{m_l}^{(l)}(k_1, k_2, \theta)}$$

## Scaling invariance for $E = 0$ :

$$f_{m_l}^{(l)}(k_1, k_2, \theta) = (k_1^2 + k_2^2)^{-(s_4+7/2)/2} (\cosh x)^{3/2} \Phi_{m_l}^{(l)}(x, \theta)$$

with  $x = \ln(k_2/k_1)$ .

The integral equation gives  $M_{s_4}^{(l)}[\vec{\Phi}^{(l)}] = 0$ .

$s_4$ allowed $\iff M_{s_4}^{(l)}$ has a zero eigenvalue
--



## RESULTS


- Numerical exploration up to  $l = 10$
- Four-body Efimov effect obtained for a single  $s_4$ , in channel  $l = 1$  with even parity:

$$D(\vec{k}_1, \vec{k}_2) = \vec{e}_z \cdot \frac{\vec{k}_1 \times \vec{k}_2}{\|\vec{k}_1 \times \vec{k}_2\|} f_0^{(1)}(k_1, k_2, \theta)$$

in the interval of mass ratio

$$\alpha_c(3; 1) \simeq 13.384 < \alpha < \alpha_c(2; 1) \simeq 13.607$$

# NUMERICAL VALUES OF $s_4 \in i\mathbb{R}$



## EXPERIMENTAL ASPECTS

- Large scattering length with magnetic Feshbach resonance (Grimm, 2006; Hulet, 2009)
- Radio-frequency spectroscopy of trimers (Jochim, 2010)
- Remaining issue: Narrow interval of mass ratio.

### Solution 1: The right mixture

- $^{41}\text{Ca}$  and  $^3\text{He}^*$  have mass ratio  $\alpha \simeq 13.58 \in [13.384, 13.607]$
- A priori,  $|s_4| \simeq 0.75$  large enough to see two tetramer states
- $^{41}\text{Ca}$  has same radioactivity as  $^{239}\text{Pu}$  (half-life  $10^5$  years)

### Solution 2: Mass tuning

- $^{40}\text{K}$  and  $^3\text{He}^*$  have slightly-off mass ratio  $\alpha \simeq 13.25$
- Use optical lattice to tune effective mass (Petrov, Shlyapnikov, 2007)

## MINLOS'S THEOREM (1995)

**Theorem:** *In the  $n + 1$  fermionic problem, the Wigner-Bethe-Peierls Hamiltonian is self-adjoint and bounded from below iff*

$$(n - 1) \frac{2\alpha(1 + 1/\alpha)^3}{\pi\sqrt{1 + 2\alpha}} \int_0^{\arcsin \frac{\alpha}{1+\alpha}} dt t \sin t < 1.$$

- We expect that “not bounded from below” is equivalent to “with Efimov effect”.
- Case  $n = 3$ :  $\alpha_c^{\text{Minlos}} \simeq 5.29$  totally differs from ours...
- Case  $\alpha = 1$ : No stable unitary gas for  $n > 9$ ...
- Weak point: Proof not included in Minlos' paper.
- Recent proof: Teta, Finco (2010). But we have found a hole in the proof. We can still **hope** that the macroscopic  $\alpha = 1$  unitary gas is stable.