

# Ab initio calculations of Be isotopes with JISP16

Pieter Maris  
pmaris@iastate.edu  
Iowa State University

IOWA STATE  
UNIVERSITY

## SciDAC project – NUCLEI

lead PIs: Joe Carlson (LANL) and Rusty Lusk (ANL)



## PetaApps award

PIs: Jerry Draayer (LSU), James P Vary (ISU),  
Ümit V Çatalyürek (OSU), Masha Sosonkina (ODU/AL)



## INCITE award – Computational Nuclear Structure

lead PI: James P Vary (ISU)



## NERSC CPU time



# Ab initio nuclear physics – Quantum many-body problem



Given a Hamiltonian operator

$$\hat{H} = \sum_{i < j} \frac{(\vec{p}_i - \vec{p}_j)^2}{2 m A} + \sum_{i < j} V_{ij} + \sum_{i < j < k} V_{ijk} + \dots$$

solve the eigenvalue problem for wave function of  $A$  nucleons

$$\hat{H} \Psi(r_1, \dots, r_A) = \lambda \Psi(r_1, \dots, r_A)$$

- eigenvalues  $\lambda$  discrete (quantized) energy levels
- eigenvectors:  $|\Psi(r_1, \dots, r_A)|^2$  probability density for finding nucleons  $1, \dots, A$  at  $r_1, \dots, r_A$



# *Ab initio nuclear physics – Computational challenges*

---

- Self-bound quantum many-body problem, with  $3A$  degrees of freedom in coordinate (or momentum) space
- Not only 2-body interactions, but also intrinsic 3-body interactions and possibly 4- and higher  $N$ -body interactions
- Strong interactions, with both short-range and long-range pieces
- Uncertainty quantification for calculations needed
  - for comparisons with experiments
  - for comparisons between different methods
- Sources of numerical uncertainty
  - statistical and round-off errors
  - systematical errors inherent to the calculational method
    - CI methods: finite basis space
    - Monte Carlo methods: sensitivity to the trial wave function
    - Lattice calculations: finite volume and lattice spacing
  - uncertainty of the nuclear potential

# Nuclear interaction


Nuclear potential not well-known ...

though in principle calculable from Quantum Chromo Dynamics

$$\hat{H} = \hat{T}_{\text{rel}} + \sum_{i < j} V_{ij} + \sum_{i < j < k} V_{ijk} + \dots$$

In practice, alphabet of realistic potentials

- Argonne potentials: AV8', AV18
  - plus Urbana 3NF (UIX)
  - plus Illinois 3NF (IL7)
- Bonn potentials
- Chiral NN interactions
  - plus chiral 3NF, ideally to the same order
- ...
- JISP16
- ...



## J-matrix Inverse Scattering Potential tuned up to $^{16}\text{O}$

- Constructed to reproduce  $np$  scattering data
- Finite rank separable potential in H.O. representation
- Nonlocal  $NN$ -only potential
- Use Phase-Equivalent Transformations (PET) to tune off-shell interaction to
  - binding energy of  $^3\text{H}$  and  $^4\text{He}$
  - low-lying states of  $^6\text{Li}$  (JISP6, precursor to JISP16)
  - binding energy of  $^{16}\text{O}$



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)



Physics Letters B 644 (2007) 33–37

PHYSICS LETTERS B

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)

## Realistic nuclear Hamiltonian: Ab exitu approach

A.M. Shirokov<sup>a,b,\*</sup>, J.P. Vary<sup>b,c,d</sup>, A.I. Mazur<sup>e</sup>, T.A. Weber<sup>b</sup>

<sup>a</sup> Skobel'syn Institute of Nuclear Physics, Moscow State University, Moscow 119992, Russia

<sup>b</sup> Department of Physics and Astronomy, Iowa State University, Ames, IA 50011-3160, USA

<sup>c</sup> Lawrence Livermore National Laboratory, L-414, 7000 East Avenue, Livermore, CA 94551, USA

<sup>d</sup> Stanford Linear Accelerator Center, MS81, 2575 Sand Hill Road, Menlo Park, CA 94025, USA

<sup>e</sup> Pacific National University, Tikhookeanskaya 136, Khabarovsk 680035, Russia

# J-matrix Inverse Scattering Potentials

---

- Constructed as matrix in H.O. basis
  - $2n + l \leq 8$  for even partial waves, limited to  $J \leq 4$
  - $2n + l \leq 9$  for odd partial waves, limited to  $J \leq 4$
  - $\hbar\omega = 40$  MeV
- $\chi^2$ /datum of 1.05 for the 1999  $np$  data base (3058 data)
- No charge symmetry breaking
- Use PET to improve
  - deuteron quadrupole moment
  - $^3\text{H}$  and  $^4\text{He}$  binding energies
  - binding energies low-lying states of  $^6\text{Li}$ : JISP6  
Shirokov, Vary, Mazur, Zaystev, Weber, PLB **621**, 96 (2005)
  - binding energy of  $^{16}\text{O}$ : JISP16  
Shirokov, Vary, Mazur, Weber, PLB **644**, 33 (2007)
  - additional tuning, more accurate calculations: JISP16<sub>2010</sub>  
reproduces  $^{16}\text{O}$  within numerical error estimates of 3%  
Shirokov, Kulikov, Maris, Mazur, Mazur, Vary, arXiv:0912.2967

# JISP16 results for few-body systems

deuteron properties

	E (MeV)	$r_p$ (fm)	$Q$ (e fm <sup>2</sup> )	$\mathcal{A}_s$ (fm <sup>-1/2</sup> )	$\mathcal{A}_d/\mathcal{A}_s$
expt.	-2.224575	1.971(6)	0.2859(3)	0.8846(9)	0.0256(4)
JISP16	-2.224575	1.964	0.2886	0.8629	0.0252
AV18	-2.224575	1.967	0.270	0.8850	0.0250

selected  $A = 3$  and 4 results

	$E_b(^3\text{H})$	$\mu(^3\text{H})$	$\mu(^3\text{He})$	$E_b(^4\text{He})$
expt.	8.482	2.979	-2.128	28.296
JISP16	8.369(2)	2.667	-1.819	28.299
AV18	7.61(1)			24.07(4)
AV18+IL2	8.43(1)	2.568(1)	-1.762(1)	28.37(3)

Pieper, Wiringa, Annu. Rev. Nucl. Part. Sci. 51, 53 (2001)

# Many-Body systems

---

## Configuration Interaction methods

- Expand wave function in basis states  $|\Psi\rangle = \sum a_i |\Phi_i\rangle$
- Express Hamiltonian in basis  $\langle \Phi_j | \hat{\mathbf{H}} | \Phi_i \rangle = H_{ij}$
- Diagonalize Hamiltonian matrix  $H_{ij}$
- Complete basis  $\rightarrow$  exact result
  - caveat: complete basis is infinite dimensional
- In practice
  - truncate basis
  - study behavior of observables as function of truncation
- Computational challenge
  - construct large  $(10^{10} \times 10^{10})$  sparse symmetric real matrix  $H_{ij}$
  - use Lanczos algorithm to obtain lowest eigenvalues & eigenvectors



# Basis space expansion

- Expand wave function in basis states  $|\Psi\rangle = \sum a_i |\Phi_i\rangle$
- Many-Body basis states  $|\Phi_i\rangle$  Slater Determinants of Single-Particle states  $|\phi\rangle$

$$\Phi_i(r_1, \dots, r_A) = \frac{1}{\sqrt{(A!)}} \begin{vmatrix} \phi_{i1}(r_1) & \phi_{i2}(r_1) & \dots & \phi_{iA}(r_1) \\ \phi_{i1}(r_2) & \phi_{i2}(r_2) & \dots & \phi_{iA}(r_2) \\ \vdots & \vdots & & \vdots \\ \phi_{i1}(r_A) & \phi_{i2}(r_A) & \dots & \phi_{iA}(r_A) \end{vmatrix}$$

- Single-Particle basis states
  - eigenstates of  $\hat{L}^2$ ,  $\hat{S}^2$ ,  $\hat{J}^2$ , and  $\hat{J}_z$  labelled by quantum numbers  $|n, l, s, j, m\rangle$
  - radial wavefunctions
    - Harmonic Oscillator
    - Wood–Saxon basis
    - Coulomb–Sturmian
    - Berggren
    - ...

Negoita, PhD thesis 2010

Caprio, Maris, Vary, PRC86, 034312 (2012)

Rotureau, last week

# Truncation scheme

---

- $M$ -scheme: Many-Body basis states eigenstates of  $\hat{\mathbf{J}}_z$

$$\hat{\mathbf{J}}_z |\Phi_i\rangle = M |\Phi_i\rangle = \sum_{k=1}^A m_{ik} |\Phi_i\rangle$$



- single run gives spectrum
- alternatives:  
 $LS$  scheme, **Coupled- $J$  scheme**, **Symplectic basis**, ...

- $N_{\max}$  truncation: Many-Body basis states satisfy

$$\sum_{k=1}^A (2n_{ik} + l_{ik}) \leq N_0 + N_{\max}$$

- exact factorization of Center-of-Mass motion
- alternatives:  
No-Core Monte-Carlo Shell Model, Importance Truncation, FCI (truncation on single-particle basis only), ...

# Intermezzo: FCI vs. $N_{\max}$ truncation



- $N_{\max}$  truncation
  - exact factorization of Center-of-Mass motion
  - converges much more rapidly than FCI truncation with basis space dimension
- Infinite basis space limit: No-Core Full Configuration (NCFC)

## Intermezzo: Center-of-Mass excitations

---

- Use single-particle coordinates, not relative (Jacobi) coordinates
  - straightforward to extend to many particles
  - have to separate Center-of-Mass motion from internal motion
- Center-of-Mass wave function **factorizes** for **H.O. basis functions** in combination with  **$N_{\max}$  truncation**

$$\begin{aligned} |\Psi_{\text{total}}\rangle &= |\phi_1\rangle \otimes \dots \otimes |\phi_A\rangle \\ &= |\Phi_{\text{Center-of-Mass}}\rangle \otimes |\Psi_{\text{int}}\rangle \end{aligned}$$

where

$$\hat{\mathbf{H}}_{\text{rel}} |\Psi_{j, \text{int}}\rangle = E_j |\Psi_{j, \text{int}}\rangle$$

- Add Lagrange multiplier to Hamiltonian (Lawson term)


$$\hat{\mathbf{H}}_{\text{rel}} \longrightarrow \hat{\mathbf{H}}_{\text{rel}} + \Lambda_{CM} \left( \hat{\mathbf{H}}_{CM}^{H.O.} - \frac{3}{2} \left( \sum_i m_i \right) \omega \right)$$

with  $\hat{\mathbf{H}}_{\text{rel}} = T_{\text{rel}} + V_{\text{rel}}$  the relative Hamiltonian



- separates CM excitations from CM ground state  $|\Phi_{CM}\rangle$

# Configuration Interaction Methods

- Expand wave function in basis states  $|\Psi\rangle = \sum a_i |\psi_i\rangle$
- Express Hamiltonian in basis  $\langle \psi_j | \hat{H} | \psi_i \rangle = H_{ij}$
- Diagonalize Hamiltonian matrix  $H_{ij}$
- **Variational**: for any finite truncation of the basis space, eigenvalue is an upper bound for the ground state energy
- **Smooth approach to asymptotic value with increasing basis space:**  
**No-Core Full Configuration** calculation
- Convergence: **independence** of  $N_{\max}$  and H.O. basis  $\hbar\omega$ 
  - different methods (NCFC, CC, GFMC, ...) using the same interaction should give same results within (statistical plus systematic) numerical uncertainties



# No-Core CI calculations – main challenge



- Increase of basis space dimension with increasing  $A$  and  $N_{\max}$
- More relevant measure for computational needs
  - number of nonzero matrix elements


# Accelerating convergence – renormalization techniques

---

Challenge: achieve numerical convergence for no-core Full Configuration calculations using finite model space calculations

- Renormalize interaction  $\longrightarrow$  effective interaction  $V_{\text{eff}}$ 
  - can improve quality of results in small model spaces
- Caveats
  - induces many-body forces
    - induced 3-body forces are often neglected
    - induced 4-, 5-, ...,  $A$ -body forces are always neglected
  - variational principle applicable to renormalized Hamiltonian not to original (bare) Hamiltonian
  - often complicates extrapolation to asymptotic values
  - need to renormalize operators as well
- Commonly used renormalization procedures
  - Lee–Suzuki effective interaction
  - Similarity Renormalization Group  
(in particular in combination with chiral interactions)


# Results with Lee–Suzuki renormalization for JISP16



- Ground state energy of  $^{16}\text{O}$  expected to be between **variational upper bound** without renormalization and **lower bound (?)** from Lee–Suzuki renormalized interaction
- Used in tuning of JISP16      Shirokov, Vary, Mazur, Weber, PLB 644, 33 (2007)



# Convergence Lee–Suzuki renormalization not monotonic



- Lee–Suzuki result for ground state energy **not** a lower bound
- JISP16 overbinds  $^{16}\text{O}$  by 10% to 15%

Maris, Vary, Shirokov, PRC79, 014308 (2009)

# Extrapolating to complete basis

---

Challenge: achieve numerical convergence for no-core Full Configuration calculations using finite model space calculations

- Perform a series of calculations with increasing  $N_{\max}$  truncation
- Extrapolate to infinite model space  $\rightarrow$  exact results
  - Empirical: binding energy exponential in  $N_{\max}$

$$E_{\text{binding}}^N = E_{\text{binding}}^{\infty} + a_1 \exp(-a_2 N_{\max})$$

- use 3 or 4 consecutive  $N_{\max}$  values to determine  $E_{\text{binding}}^{\infty}$
- use  $\hbar\omega$  and  $N_{\max}$  dependence to estimate numerical error bars



Maris, Shirokov, Vary, PRC79, 014308 (2009)

- Recent studies of IR and UV behavior
  - exponentials in  $\sqrt{\hbar\omega/N}$  and  $\sqrt{\hbar\omega N}$  Coon *et al*, arXiv:1205.3230; Furnstahl, Hagen, Papenbrock PRC86, 031301(R) (2012)

# Extrapolating to complete basis – in practice

- Perform a series of calculations with increasing  $N_{\max}$  truncation
- Use empirical exponential in  $N_{\max}$ :

$$E_{\text{binding}}^N = E_{\text{binding}}^{\infty} + a_1 \exp(-a_2 N_{\max})$$



- H.O. basis up to  $N_{\max} = 16$ :  $E_b = -31.49(3)$  MeV

Cockrell, Maris, Vary, PRC86 034325 (2012)


- Hyperspherical harmonics up to  $K_{\max} = 14$ :  $E_b = -31.46(5)$  MeV

Vaintraub, Barnea, Gazit, PRC79 065501 (2009)


# Accelerating convergence – Coulomb-Sturmian basis

Caprio, Maris, Vary, PRC86, 034312 (2012)

- Asymtotic behavior
  - H.O. basis  $\exp(-a r^2)$
  - Coulomb–Sturmian basis  $\exp(-c r)$
- Disadvantage
  - no exact factorization of Center-of-Mass motion
  - in practice, approximate factorization  
Hagen, Papenbrock, Dean, PRL103, 062503 (2009)
  - can use Lagrange multiplier to remove spurious state



# Coulomb-Sturmian – binding energies



● at  $N_{\text{max}} = 4$  further from convergence than H.O. basis


● extrapolate to the same results as H.O. basis

● dashed line: extrapolated result from  $N_{\text{max}} = 16$  calculations in H.O. basis

Cockrell, Maris, Vary,  
PRC86 034325 (2012)

# Coulomb-Sturmian – radius

Caprio, Maris, Vary, PRC86, 034312 (2012)




- exponential extrapolation does not work for radii in H.O. basis
- exponential extrapolation seems to work for radii in C.S. basis
- best estimate based on  $N_{\text{max}} = 16$  H.O. calculations: 2.3 fm



Cockrell, Maris, Vary, PRC86 034325 (2012)

- experimental point-proton radius: 2.45 fm

# Ground state energy Be-isotopes with JISP16



# *7*Be – Ground state properties





- Binding energy converges monotonically, with optimal H.O. frequency around  $\hbar\omega = 20$  MeV to 25 MeV
- Ground state about 0.7 MeV underbound with JISP16
- Proton point radius does not converge monotonically
  - Coulomb–Sturmian basis likely to improve convergence




# 7Be – Proton density

- Translationally-invariant density – center-of-mass motion taken out  
w. Cockrell, PhD thesis 2012




- Slow build up of asymptotic tail of wavefunction
- Proton density appears to converge more rapidly at  $\hbar\omega = 12.5$  MeV than at 20 MeV because long-range part of wavefunction is better represented with smaller H.O. parameter

# 7Be – Proton radius





- Calculation one-body observables  $\langle i | \mathcal{O} | j \rangle \sim \int \mathcal{O}(r) r^2 \rho_{ij}(r) dr$
- RMS radius:  $\mathcal{O}(r) = r^2$
- Slow convergence of RMS radius due to slow build up of asymptotic tail
- Ground state RMS radius in agreement with data

# *7*Be – Quadrupole moment



- Ground state quadrupole moment in agreement with data
- Optimal basis space around  $\hbar\omega = 10$  MeV to 12 MeV
- Similar slow convergence for E2 transitions

# 7Be – Excited states





- Excitation energy of narrow states
  - converge rapidly
  - agree with experiments
- Broad resonances depend  $\hbar\omega$

- Magnetic moments well converged
  - 2-body currents needed for agreement with data (meson-exchange currents)

# 7Be – Emergence of rotational band? *in progress, w. M. Caprio*

E2 observables suggest rotational structure for  $\frac{3}{2}^-$ ,  $\frac{1}{2}^-$ ,  $\frac{7}{2}^-$ ,  $\frac{5}{2}^-$  states




$$Q(J) = \frac{\frac{3}{4} - J(J+1)}{(J+1)(2J+3)} Q_0$$

$$B(E2; i \rightarrow f) = \frac{5}{16\pi} Q_0^2 \left( J_i, \frac{1}{2}; 2, 0 \middle| J_f, \frac{1}{2} \right)^2$$



# 7Be – Structure of $(\frac{5}{2}^-, \frac{1}{2})_1$ (broad) and $(\frac{5}{2}^-, \frac{1}{2})_2$ (narrow) states

Translationally-invariant nucleon densities


Cockrell, PhD thesis 2012





proton  
density



neutron  
density




# 9Be – Ground state properties



- Convergence pattern natural and unnatural parity looks similar
- Ground state about  $1.0 \pm 0.2$  MeV underbound with JISP16
- Lowest unnatural parity state underbound by about  $2.7 \pm 0.8$  MeV
  - need next basis space for unnatural parity
  - need improved interaction?



# Positive vs. negative parity states of Be-isotopes



- Unnatural parity states systematically underbound by about 1 MeV to 2 MeV compared to lowest natural parity states
  - interaction JISP16 not good enough?
  - difference in convergence of pos. and neg. parity states?





# 9Be – Positive and negative spectrum



- Excitation energy  $5/2^-$  at 3 MeV well converged (narrow)
- Excitation energy  $7/2^-$  reasonably converged
- Excitation energies broad neg. parity not well converged
- Excitation energies pos. parity well converged

# 9Be – Emergence of rotational bands *in progress, w. M. Caprio*





- Rotational energy for states with axial symmetry  $E(J) \propto J(J + 1)$
- Quadrupole moments for rotational band


$$Q(J) = \frac{3K^2 - J(J + 1)}{(J + 1)(2J + 3)} Q_0$$

- Quadrupole moments not converged, but ratio of quadrupole moments agree with rotational band structure

# $9\text{Be}$ – Structure: Density ( $\frac{3}{2}^-, \frac{1}{2}$ ) ground state





and their difference



Translationally-invariant  
proton and neutron densities  
Chase Cockrell, PhD thesis, 2012



● Emergence of  $\alpha$  clustering?

# Rotational bands odd Be isotopes *in preparation, w. M. Caprio*





- Quadrupole moments not converged, but ratio of quadrupole moments agree with rotational band structure

- Also for the unnatural parity states




- Quadrupole moments not converged, but ratio of quadrupole moments agree with rotational band structure


# Rotational bands even Be isotopes *in preparation, w. M. Caprio*



# $B(E2)$ transitions Be isotopes *in preparation, w. M. Caprio*



- Rotor prediction
- $\Delta_J = 1$ : dashed
- $\Delta_J = 2$ : solid



● Ratio's  $B(E2)/Q^2$  in agreement with rotational structure as well

# Conclusions

---

- **No-core Configuration Interaction** nuclear structure calculations
  - **Binding energy**, spectrum
  - $\langle r^2 \rangle$ ,  $\mu$ ,  $Q$ , transitions, wfns, one-body densities
- Main challenge: construction and diagonalization of extremely large ( $D > 1$  billion) sparse matrices
- Need realistic basis function to improve convergence  $\langle r^2 \rangle$ ,  $Q$
- **JISP16**
  - Nonlocal phenomenological 2-body interaction
  - Good description of a range of light nuclei
  - Rapid convergence for binding energies
  - Emergence of rotational bands and clustering in Be-isotopes
- Would not have been possible without collaboration with applied mathematicians and computer scientists  
Aktulga, Yang, Ng (LBNL); Çatalyürek, Saule (OSU); Sosonkina (ODU/AL)