

# Energy Loss at “NLO” in eXtremely hot QCD

Derek Teaney

SUNY Stony Brook and RBRC Fellow



Stony Brook University

- Jacopo Ghiglieri, J. Hong, A. Kurkela, G. Moore, DT, JHEP
- [Jacopo Ghiglieri](#), G. Moore, DT, arXiv:almost-done
- Jacopo Ghiglieri, G. Moore, DT, B. Schenke, just-starting

## Outline: Energy loss and transport in weakly coupled plasmas at “NLO”


1. Philosophy of weakly coupled calculations – there is only one right answer . . .
  - (a) Collisional vs. radiative loss
  - (b) Corrections to collinear formalism
  - (c) Relation between drag and radiative loss
  - (d) Can work with enough kicking
2. Energy loss of light quarks and gluons
3. Thermal photons

Jacopo Ghiglieri, G. Moore, DT


Jacopo Ghiglieri, J. Hong, A. Kurkela, G. Moore, DT, JHEP

## Three mechanisms for energy loss and transport at LO in QGP

### 1. Hard Collisions: $2 \leftrightarrow 2$




### 2. Drag: collisions with soft random classical field



$$\frac{dp}{dt} = -\eta \hat{v}$$

### 3. Brem: $1 \leftrightarrow 2$

- random walk induces collinear bremsstrahlung



- The probability of a transverse kick of momentum  $\mathbf{q}_\perp$  from soft fields:

$$\hat{C}_{LO}[\mathbf{q}_\perp] = \frac{Tm_D^2}{q_\perp^2 (q_\perp^2 + m_D^2)}$$


NLO involves corrections to these processes and the relation between them.

Same processes determine the shear viscosity of QCD in high temperature plasma!



Three rates for energy loss at leading order:


1. Hard Collisions – a  $2 \leftrightarrow 2$  processes



$$[\partial_t + v_{\mathbf{k}} \cdot \partial_{\mathbf{x}}] f_{\mathbf{k}} = C_{2 \leftrightarrow 2}[\mu_{\perp}]$$

Total  $2 \leftrightarrow 2$  scattering rate depends logarithmically on the cutoff

## 2. Drag and long-diffusion: A longitudinal force-force correlator along the light cone



$$[\partial_t + v_{\mathbf{k}} \cdot \partial_{\mathbf{x}}] f_{\mathbf{k}} = \eta(\mu) \mathbf{v} \cdot \frac{\partial f_{\mathbf{k}}}{\partial \mathbf{k}}$$

- Evaluate longitudinal force-force with hard thermal loops + sum-rules


$$\eta(\mu) \propto g^2 C_A \int^{\mu} \frac{d^2 \mathbf{p}_T}{(2\pi)^2} \int \frac{dp_+ dp_0}{(2\pi)^4} \underbrace{\langle F_{z+}(P) F_{z+} \rangle 2\pi \delta(p_+)}_{\text{evaluate with sum-rule } p_0 \rightarrow \infty}$$

$$\propto g^2 C_A \int^{\mu} \frac{d^2 \mathbf{p}_T}{(2\pi)^2} \frac{m_{\infty}^2}{p_T^2 + m_{\infty}^2}$$

$$\propto g^2 C_A \frac{m_{\infty}^2}{4\pi} \log(\mu^2 / m_{\infty}^2)$$

The  $\mu$ -dependence of the drag cancels against  $\mu$ -dependence of the  $2 \rightarrow 2$  rate

### 3. Collinear Bremsstrahlung – a $1 \leftrightarrow 2$ processes



$$[\partial_t + v_{\mathbf{k}} \cdot \partial_x] f_{\mathbf{k}} =$$

$$\underbrace{C_{1 \leftrightarrow 2}}$$

LPM + AMY and all that stuff!

The bremsstrahlung rate is proportional to the rate of transverse momentum kicks,  $\hat{C}_{LO}[\mathbf{q}_{\perp}]$ :

$$\hat{C}_{LO}[q_{\perp}] = \text{in medium scattering rate with momentum } \mathbf{q}_{\perp}$$

- Need to compute transverse force-force correlators along the light cone.

$$q_{\perp}^2 \hat{C}_{LO}[\mathbf{q}_{\perp}] = \int \frac{dq_+ dq_0}{(2\pi)^2} \underbrace{\langle F_{i+}(Q) F_{i+} \rangle}_{\text{evaluate with sum rule at } q_0 = 0} 2\pi \delta(q_+)$$

$$= \frac{T m_D^2}{q_{\perp}^2 + m_D^2}$$



Summary – the full LO Boltzmann equation:

$$[\partial_t + v_{\mathbf{k}} \cdot \partial_{\mathbf{x}}] f_{\mathbf{k}} = \eta(\mu) \mathbf{v}_{\mathbf{k}} \cdot \frac{\partial f_{\mathbf{k}}}{\partial \mathbf{k}} + C_{2 \leftrightarrow 2}[\mu] + C_{1 \leftrightarrow 2}$$

$\mu$  dependence cancels

The diagram features a red horizontal line with two downward-pointing arrows. The left arrow points to the  $\eta(\mu)$  term in the equation, and the right arrow points to the  $C_{2 \leftrightarrow 2}[\mu]$  term. This indicates that the  $\mu$  dependence of the drag term cancels with the  $\mu$  dependence of the  $2 \rightarrow 2$  rate.


soft sector drag      soft sector momentum broadening  $\hat{C}_{LO}[q_{\perp}]$

1) The cutoff dependence of the drag cancels against the  $2 \rightarrow 2$  rate!

2) Soft sector enters in just a few places.

3) Light cone sum rules.

## Hot QGP with a Jet




$$\overline{\frac{dE}{dx}} = \text{Energy loss rate}$$

Use the Boltzmann equation for energy loss or shear viscosity:

$$\overline{\frac{dE}{dx}} \propto g^2 T^2 \left[ \underbrace{O(g^2 \log) + O(g^2)}_{\text{LO Boltzmann (AMY)}} + \underbrace{O(g^3 \log) + O(g^3)}_{\text{NLO, from soft } gT \text{ gluons, } n_B \simeq \frac{T}{\omega} \simeq \frac{1}{g}} \right] + \dots$$

## $O(g)$ Corrections to Hard Collisions, Drag, Bremm:

1. No corrections to Hard Collisions:
2. Corrections to Drag:




- Nonlinear interactions of soft classical fields changes the force-force correlator
- Doable because of HTL sum rules (light cone causality)

### 3. Corrections to Breemm:

(a) Small angle breemm. Corrections to AMY coll. kernel.


(Caron-Huot)



$$\hat{C}_{LO}[q_\perp] = \frac{T g^2 m_D^2}{q_\perp^2 (q_\perp^2 + m_D^2)} \rightarrow \text{A complicated but analytic formula}$$

(b) Large angle brem and collisions with plasmons.


- Include collisions with energy exchange,  $q^- \sim gT$ .



The large-angle (semi-collinear radiation) interpolates collisional and rad. loss

The NLO Boltzmann equation – a preview:


Cutoff dependence cancels

$$[\partial_t + v_{\mathbf{k}} \cdot \partial_{\mathbf{x}}] f_{\mathbf{k}} = (\eta(\mu) + \delta\eta(\mu)) \mathbf{v}_{\mathbf{k}} \cdot \frac{\partial f_{\mathbf{k}}}{\partial \mathbf{k}} + C_{2\leftrightarrow 2}[\mu]$$
$$C_{1\leftrightarrow 2} + \delta C_{1\leftrightarrow 2} + C_{\text{semi-coll}}[\mu]$$


The  $\mu$ -dependence of the drag at NLO cancels the  $\mu$ -dependence of semi-collinear radiation

## Semi-collinear radiation – a new kinematic window


$2 \rightarrow 2$  processes




The semi-collinear regime interpolates between brem and collisions

## Matching collisions to brem

- When the gluon becomes soft (a plasmon), the  $2 \leftrightarrow 2$  collision:



is *not* physically distinct from the wide angle brem




Need both processes

- For harder gluons,  $q^- \rightarrow T$ , bremm becomes a normal  $2 \rightarrow 2$  process.
- For softer gluons,  $q^- \rightarrow g^2 T$ , wide angle bremm matches onto collinear limit.

## Brem and collisions at wider angles (but still small!)

- Semi-collinear emission:



- The matrix element is:

$$|\mathcal{M}|^2 (2\pi)^4 \delta^4(P_{\text{tot}}) \propto \underbrace{\frac{1+z^2}{z}}_{\text{QCD splitting fcn}} \int_Q \frac{1}{(q^-)^2} \underbrace{\langle F_{i+}(Q) F_{i+} \rangle}_{\text{scattering-center}} 2\pi \delta(q^- - \delta E)$$


All of the dynamics of the scattering center in a single matrix element  $\langle F_{i+}(Q) F_{i+} \rangle$




The scattering center:

$$\hat{C}[\mathbf{q}_\perp, \delta E] = \int_Q \frac{1}{(q^-)^2} \langle F_{i+}(Q) F_{i+} \rangle 2\pi \delta(q^- - \delta E)$$

1. Soft-correlator has wide angle brem = cut
2. And plasmon scattering = poles



## Finite energy transfer sum-rule



- The small angle brems rate involves:

$$\underbrace{q_{\perp}^2 \hat{C}_{LO}[q_{\perp}] = \int_{-\infty}^{\infty} \frac{dq_0}{2\pi} \langle F_{i+} F_{i+}(Q) \rangle |_{q_+=0} = \frac{T m_D^2}{q_T^2 + m_D^2}}_{\text{Rate of transverse kicks of } q_{\perp}}$$

- The wide angle brems rate involves a finite  $q^- = \delta E$  generalization:


$$\underbrace{\int_{-\infty}^{\infty} \frac{dq_0}{2\pi} \langle F_{i+} F_{i+}(Q) \rangle |_{q_+=-\delta E} = T \left[ \frac{2(\delta E)^2 (\delta E^2 + q_{\perp}^2 + m_D^2) + m_D^2 q_{\perp}^2}{(\delta E^2 + q_{\perp}^2 + m_D^2)(\delta E^2 + q_{\perp}^2)} \right]}_{\text{Rate of transverse kicks of } q_{\perp} \text{ and energy transfer } \delta E}$$

almost involves the replacement,  $q_{\perp}^2 \rightarrow q_{\perp}^2 + \delta E^2$



## Matching between brem and drag

$2 \rightarrow 2$  processes




- The semi-collinear emission rate diverges logarithmically when the gluon gets soft

$$\Gamma_{\text{semi-coll}} \sim g^2 C_A \overbrace{\frac{\delta m_\infty^2}{4\pi}}{\sim g^3 T^2} \log \left( \frac{2T m_D}{\mu} \right)$$


When the gluon becomes soft need to relate radiation and drag.

## Matching between semi-collinear brem and drag

- When the final gluon line becomes soft, the brem process:



is *not* physically distinct from the drag process:




but represents a higher order correction to drag.

Separately both processes depend on the separation scale,  $\mu \sim gT$ , but . . .

the  $\mu$  dep. cancels when both rates are included

## Computing the NLO drag:



- Evaluate NLO longitudinal force-force with hard thermal loops + sum-rules
- Only change relative to LO is the replacement  $m_\infty^2 \rightarrow m_\infty^2 + \delta m_\infty^2$

$$\eta(\mu) \propto g^2 C_A \int^\mu \frac{d^2 \mathbf{p}_T}{(2\pi)^2} \frac{m_\infty^2 + \delta m_\infty^2}{p_T^2 + m_\infty^2 + \delta m_\infty^2}$$

$$\propto \text{leading order} + \underbrace{g^2 C_A \frac{\delta m_\infty^2}{4\pi} \left[ \log \left( \frac{\mu_\perp^2}{m_\infty^2} \right) - 1 \right]}_{\text{NLO correction to drag}}$$

The cutoff dependence of the drag cancels against the semi-collinear emission rate

## The NLO Boltzmann equation review:

Cutoff dependence cancels

$$[\partial_t + v_{\mathbf{k}} \cdot \partial_{\mathbf{x}}] f_{\mathbf{k}} = (\eta(\mu) + \delta\eta(\mu)) \mathbf{v}_{\mathbf{k}} \cdot \frac{\partial f_{\mathbf{k}}}{\partial \mathbf{k}} + C_{2\leftrightarrow 2}[\mu]$$
$$C_{1\leftrightarrow 2} + \delta C_{1\leftrightarrow 2} + C_{\text{semi-coll}}[\mu]$$

## Lessons from weak coupling


- Tight relation between drag, wide angle emissions, quasi-particle mass shift.
  - Closely related to dimensional reduction.
- The wide angle emission kernel  $\hat{C}[\mathbf{q}_{\perp}, \delta E]$  is closely related to  $\hat{C}[\mathbf{q}_{\perp}]$ , almost:

$$\mathbf{q}_{\perp}^2 \rightarrow \mathbf{q}_{\perp}^2 + \delta E^2$$

- Closely related to dimensional reduction.
- Understand in detail the transition from radiative to collisional loss

Currently being implemented into e-loss models, e.g. MARTINI

## Simulation of a 20 GeV gluon







$$2k(2\pi)^3 \frac{d\Gamma}{d^3k} = \text{Photon emission rate per phase-space}$$



Same techniques can be used for thermal photon production:

- The rate is function of the coupling constant and  $k/T$ :

$$2k(2\pi)^3 \frac{d\Gamma}{d^3k} \propto e^2 T^2 \left[ \underbrace{O(g^2 \log) + O(g^2)}_{\text{LO AMY}} + \underbrace{O(g^3 \log) + O(g^3)}_{\text{From soft } gT \text{ gluons, } n_B \simeq \frac{T}{\omega} \simeq \frac{1}{g}} \right] + \dots$$



$O(g^3)$  is closely related to  $O(g^3)$  in energy loss:

## NLO Results:



NLO corrections are modest and roughly  $k$  independent

The different contributions at NLO (conversions are not numerically important)



## Conclusion:

- NLO corrections to collinear processes seem to be modest.
- All of the soft sector buried into a few coefficients, e.g.  $\hat{C}[q_{\perp}, \delta E]$  and  $\delta m_{\infty}^2$ 
  - Can we compute these non-perturbatively with dimensional reduction?
    - \* First start: Marco Panero, Kari Rummukainen, Andreas Schafer arxiv:1307.5850
  - Use these non-perturbative parameters to compute  $\eta/s$

A program for computing QGP transport perturbatively with non-perturbative inputs for the Debye and magnetic sector