

# Nonlocal Dispersive Optical Model

Hossein Mahzoon  
Wim. Dickhoff  
Robert Charity  
Helber Dussan  
Seth Waldecker

**Washington University in St. Louis**  
March 2015

# Table of contents

- Introduction to DOM
- Role of Nonlocality
- Results for  $^{40}\text{Ca}$  and  $^{48}\text{Ca}$
- Spectral Functions

# Hartree-Fock Potential

In general we want to solve a problem with Hamiltonian

$$H = T + V$$

The irreducible self energy can be written as:

$$\Sigma^*(y, x; E) = -i \int \frac{dE'}{2\pi} \sum_{x', y'} \langle yx' | V | xy' \rangle G(y', x'; E') + \text{Higher Orders}(E)$$

The Hartree-Fock approximation means to eliminate “higher orders” which in general depend on energy.

# Dispersion Relation



By evaluating the real part, let say at some energy  $\varepsilon_F$  , one can rewrite the dispersion relation as:

$$\text{Re}\Sigma(x, y; E) = \Sigma_s(x, y) - \mathcal{P} \int_{\varepsilon_T^+}^{\infty} \frac{dE'}{\pi} \frac{\text{Im}\Sigma(x, y; E')}{E - E'} + \mathcal{P} \int_{-\infty}^{\varepsilon_T^-} \frac{dE'}{\pi} \frac{\text{Im}\Sigma(x, y; E')}{E - E'}$$

$$\text{Re}\Sigma(x, y; \varepsilon_F) = \Sigma_s(x, y) - \mathcal{P} \int_{\varepsilon_T^+}^{\infty} \frac{dE'}{\pi} \frac{\text{Im}\Sigma(x, y; E')}{\varepsilon_F - E'} + \mathcal{P} \int_{-\infty}^{\varepsilon_T^-} \frac{dE'}{\pi} \frac{\text{Im}\Sigma(x, y; E')}{\varepsilon_F - E'}$$

$$\begin{aligned} \text{Re}\Sigma(x, y; E) = \text{Re}\Sigma(x, y; \varepsilon_F) - \mathcal{P} \int_{\varepsilon_T^+}^{\infty} \frac{dE'}{\pi} \text{Im}\Sigma(x, y; E') \times \left[ \frac{1}{E - E'} - \frac{1}{\varepsilon_F - E'} \right] \\ + \mathcal{P} \int_{-\infty}^{\varepsilon_T^-} \frac{dE'}{\pi} \text{Im}\Sigma(x, y; E') \times \left[ \frac{1}{E - E'} - \frac{1}{\varepsilon_F - E'} \right] \end{aligned}$$

# Effect of dispersion relation



*C. Mahaux and R. Sartor, Adv. Nucl. Phys.*

# Lehmann representation:

$$G(\alpha, \beta; t - t') = -\frac{i}{\hbar} \langle \Psi_0^N | \mathcal{T} [a_{\alpha_H}(t) a_{\beta_H}^\dagger(t')] | \Psi_0^N \rangle$$

$$G(\alpha, \beta; E) = \sum_m \frac{\langle \Psi_0^N | a_\alpha | \Psi_m^{N+1} \rangle \langle \Psi_m^{N+1} | a_\beta^\dagger | \Psi_0^N \rangle}{E - (E_m^{N+1} - E_0^N) + i\eta} + \sum_n \frac{\langle \Psi_0^N | a_\beta^\dagger | \Psi_n^{N-1} \rangle \langle \Psi_n^{N-1} | a_\alpha | \Psi_0^N \rangle}{E - (E_0^N - E_n^{N-1}) - i\eta}$$
$$= \langle \Psi_0^N | a_\alpha \frac{1}{E - (\hat{H} - E_0^N) + i\eta} a_\beta^\dagger | \Psi_0^N \rangle + \langle \Psi_0^N | a_\beta^\dagger \frac{1}{E - (E_0^N - \hat{H}) - i\eta} a_\alpha | \Psi_0^N \rangle$$

$$G(\alpha, \beta; E) = G^{(0)}(\alpha, \beta; E) + \sum_{\gamma, \delta} G(\alpha, \gamma; E) \Sigma^*(\gamma, \delta; E) G^{(0)}(\delta, \beta; E)$$

Dyson  
equation

# Local DOM Potential

$$U = \mathcal{V} + i\mathcal{W}$$

$$\mathcal{W}(r, E) = -\mathcal{W}_v(r, E)f(r, r_v, a_v) + 4a_s W_s(E) \frac{d}{dr} f(r, r_s, a_s) + \mathcal{W}_{so}(r, E)$$

$$\mathcal{V}(r, E) = \mathcal{V}_{HF}(r, E) + \Delta\mathcal{V}(r, E)$$

$$\Delta\mathcal{V}(r, E) = \frac{1}{\pi} \mathcal{P} \int \mathcal{W}(r, E') \left( \frac{1}{E' - E} - \frac{1}{E' - \varepsilon_F} \right) dE'$$

$$\mathcal{V}_{HF}(r, E) = -V_{HF}^{Vol}(E)f(r, r_{HF}, a_{HF}) + 4V_{HF}^{Sur} \frac{d}{dr} f(r, r_{HF}, a_{HF}) + V_c(r) + \mathcal{V}_{so}(r, E)$$

Where

$$f(r, r_i, a_i) = \frac{1}{1 + e^{\frac{r-r_i A^{1/3}}{a_i}}}$$

## Nonlocal extension of the dispersive optical model to describe data below the Fermi energy

W. H. Dickhoff,<sup>1</sup> D. Van Neck,<sup>2</sup> S. J. Waldecker,<sup>1</sup> R. J. Charity,<sup>3</sup> and L. G. Sobotka<sup>1,3</sup>

<sup>1</sup>*Department of Physics, Washington University, St. Louis, Missouri 63130, USA*


<sup>2</sup>*Center for Molecular Modeling, Ghent University, Technologiepark 903, B-9052 Zwijnaarde, Belgium*

<sup>3</sup>*Department of Chemistry, Washington University, St. Louis, Missouri 63130, USA*

(Received 1 October 2010; published 10 November 2010)

$$\sum_{HF} (r, r') = V_{NL} f\left(\frac{1}{2}|r + r'|\right) H(|r - r'|)$$


$$H(|r - r'|) = \frac{1}{\pi^{\frac{3}{2}} \beta^3} \exp\left[-\left(\frac{r - r'}{\beta}\right)^2\right]$$






# Role of Nonlocality

CDBonn (short range)



*Phys. Rev. C 84, 044319 (2011)*

FRPA



*Phys. Rev. C 84, 034616 (2011)*

# Nonlocal DOM


- For Example:

$$\text{Im } \Sigma(r \llcorner r' \llcorner E) = \text{Im } \Sigma^{\text{nl}}(r \llcorner r'; E) + \delta(r - r') \mathcal{W}^{\text{so}}(r; E)$$



$$\text{Im } \Sigma^{\text{nl}}(r \llcorner r'; E) = -W_{0\pm}^{\text{vol}}(E) f(\tilde{r}; r_{\pm}^{\text{vol}}; a_{\pm}^{\text{vol}}) H(\mathbf{s}; \beta_{\text{vol}}^{\pm})$$

$$+ 4a_{\pm}^{\text{sur}} W_{\pm}^{\text{sur}}(E) H(\mathbf{s}; \beta_{\text{sur}}^{\pm}) \frac{d}{d\tilde{r}} f(\tilde{r} \llcorner r_{\pm}^{\text{sur}} \llcorner a^{\text{sur}})$$


# $^{40}\text{Ca}$ Cross section



# $^{40}\text{Ca}$ Cross sections and analyzing power



# The hole spectral function for high momenta




*Data:(dotted-line)*  
 D. Rohe, Habilitationsschrift  
 (University of Basel,  
 Basel,2004)

*Nonlocal-DOM:(dashed-  
 dotted)*

$$S_h(E_m, p_m) = \sum_n \delta(E_m - E_0^N - E_n^{N-1}) |\langle \Psi_n^{N-1} | a_{p_m} | \Psi_0^N \rangle|^2$$


# $^{40}\text{Ca}$ Charge Density

Local DOM



PRC **82**, 054306(2010)

NonLocal DOM




PRL **112**, 162503(2014)

# $^{48}\text{Ca}$ Cross section


- Including Asymmetry terms proportional to  $\frac{N - Z}{A}$
- All the parameters kept fixed except the radii (comparing to  $^{40}\text{Ca}$ )

# $^{48}\text{Ca}$ Cross sections

Nonlocal




local







# $^{48}\text{Ca}$ Cross sections



PRC **83**, 064605 (2011)



PRL **112**, 162503(2014)



# Spectroscopic Factors

protons

	<b>40Ca</b>	<b>48Ca</b>
1s12	0.73	0.63
0d32	0.76	0.69
0f72	0.73	0.63



neutrons

	<b>40Ca</b>	<b>48Ca</b>
1s12	0.76	0.80
0d32	0.78	0.77
0f72	0.71	0.80

# Weak charge

- The electron interacts with the nucleus by exchanging either a photon or  $Z^0$  boson.
- $Z^0$  boson has a much larger coupling to the neutron than protons.

# $^{48}\text{Ca}$ Charge Density



Eur. Phys. J. A (2014) 50:  
48 J. Horowitz, K.S. Kumar, and R. Michaels

# Spectral Function

$$S_{\ell j}^p(k, k'; E) = \frac{i}{2\pi} \left[ G_{\ell j}^p(k, k'; E^+) - G_{\ell j}^p(k, k'; E^-) \right]$$

$$G_{\ell j}^p(k, k'; E^\pm) = \sum_n \frac{\phi_{\ell j}^{n+}(k) \left[ \phi_{\ell j}^{n+}(k') \right]^*}{E - E_n^{*A+1} \pm i\eta} + \sum_c \int_{T_c}^\infty dE' \frac{\chi_{\ell j}^{cE'}(k) \left[ \chi_{\ell j}^{cE'}(k') \right]^*}{E - E' \pm i\eta}$$

$$\phi_{\ell j}^{n+}(k) = \langle \Psi_0^A | a_{k\ell j} | \Psi_n^{A+1} \rangle$$


$$\chi_{\ell j}^{cE}(k) = \langle \Psi_0^A | a_{k\ell j} | \Psi_{cE}^{A+1} \rangle$$

# Spectral Function

$$S_{\ell j}^p(r, r'; E) = \sum_c \chi_{\ell j}^{cE}(r) [\chi_{\ell j}^{cE}(r')]^* \quad \frac{k^2}{2m} \phi_{\ell j}^n(k) + \int dq q^2 \operatorname{Re} \Sigma_{\ell j}^*(k, q; \varepsilon_n) \phi_{\ell j}^n(q) = \varepsilon_n \phi_{\ell j}^n(k)$$

$$S_{\ell j}^{n-}(E) = \int dr r^2 \int dr' r'^2 \phi_{\ell j}^{n-}(r) S_{\ell j}^h(r, r'; E) \phi_{\ell j}^{n-}(r'),$$



$$S_{\ell j}^{n+}(E) = \int dr r^2 \int dr' r'^2 \phi_{\ell j}^{n-}(r) S_{\ell j}^p(r, r'; E) \phi_{\ell j}^{n-}(r'),$$



**In Practice**  $\rightarrow S_{\ell j}^p(k, k'; E) = \frac{i}{2\pi} \left[ G_{\ell j}^p(k, k'; E^+) - G_{\ell j}^p(k, k'; E^-) \right]$

$$S_{\ell j}^p(r, r'; E) = \frac{2}{\pi} \int dk k^2 \int dk' k'^2 j_\ell(kr) S_{\ell j}^p(k, k'; E) j_\ell(k'r'),$$

# Spectral Strength




orbit	$n_{nlj}$	$d_{nlj}[0, 200]$	$n_{nlj} + d_{nlj}[\epsilon_F, 200]$	$d_{nlj}[0, 200]$
	DOM	DOM	DOM	CDBonn
$0s_{1/2}$	0.926	0.032	0.958	0.035
$0p_{3/2}$	0.914	0.047	0.961	0.036
$0p_{1/2}$	0.906	0.051	0.957	0.038
$0d_{5/2}$	0.883	0.081	0.964	0.040
$1s_{1/2}$	0.871	0.091	0.962	0.038
$0d_{3/2}$	0.859	0.097	0.966	0.041
$0f_{7/2}$	0.046	0.202	0.970	0.034
$0f_{5/2}$	0.036	0.320	0.947	0.036


# Spectral Function

$$\chi_{lj}^{cE}(k) = \langle \Psi_0^A | a_{k\ell j} | \Psi_{cE}^{A+1} \rangle \quad \chi_{lj}^{elE}(r) = \left[ \frac{2mk_0}{\pi\hbar^2} \right]^{1/2} \left\{ j_\ell(k_0 r) + \int dk k^2 j_\ell(kr) G^{(0)}(k; E) \Sigma_{\ell j}(k, k_0; E) \right\}$$

$l = 4$



$l = 2$







# Spectral Function

- Screened Coulomb

$$w_R(r) = w(r)e^{-(r/R)^n}$$

Phys. Rev. C 41, 2615 (1990)



0s12	0p12	0p32	0f72
0.94	0.93	0.95	0.95

← sum rule

# Conclusion :

- According to the results, Nonlocal DOM is a reliable candidate to study nuclear properties.
- Nonlocality and Dispersion corrections playing an important role to get the physics of the system correctly.

