

Physics 570 — Handout on Dimensional Regularization

Consider the following integral in n dimensions with a *Euclidian* metric:

$$I_1 \equiv \int d^n k \frac{1}{(k^2 + a^2)^r} .$$

We may evaluate this making in terms of the Γ function:

$$\alpha^{-s} \Gamma(s) = \int_0^\infty dx x^{s-1} e^{-\alpha x} .$$

Then

$$\begin{aligned} I_1 &= \frac{1}{\Gamma(r)} \int d^n k \int_0^\infty dx x^{r-1} e^{-x(k^2 + a^2)} \\ &= \frac{\pi^{n/2}}{\Gamma(r)} \int_0^\infty dx x^{r-1-n/2} e^{-xa^2} \\ &= \pi^{n/2} a^{n-2r} \frac{\Gamma(r - n/2)}{\Gamma(r)} \end{aligned}$$

Another useful integral is

$$I_2 \equiv \int d^n k \frac{k^2}{(k^2 + a^2)^r} .$$

To get this we define

$$\begin{aligned} I_1(\alpha) &\equiv \int d^n k \frac{1}{(\alpha k^2 + a^2)^r} ; \\ &= \alpha^{-n/2} I_1 \end{aligned}$$

then by differentiating by α and setting $\alpha = 1$ we find

$$I_2 = \frac{n\pi^{n/2} a^{n-2r+2}}{2(r-1)} \frac{\Gamma(r-1-n/2)}{\Gamma(r-1)} .$$

Finally note that

$$\begin{aligned} I_3^{\mu\nu} &\equiv \int d^n k \frac{k^\mu k^\nu}{(k^2 + a^2)^r} . \\ &= \frac{\delta^{\mu\nu}}{n} I_2 \end{aligned}$$

Useful Properties of Γ Functions

Gamma functions have the property $\Gamma(z+1) = z\Gamma(z)$, with $\Gamma(1) = 1$. Thus for integers $n \geq 1$,

$$\Gamma(n+1) = n!, \quad n \geq 1.$$

Also useful is the value

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$

The Gamma function is singular for non-positive integer arguments. Near these singularities it can be expanded as

$$\Gamma(-n + \epsilon) = \frac{(-1)^n}{n} \left[\frac{1}{\epsilon} + \psi(n+1) + \mathcal{O}(\epsilon) \right],$$

where

$$\begin{aligned} \psi(n+1) &= 1 + \frac{1}{2} + \dots + \frac{1}{n} - \gamma, \\ \gamma &= 0.5772\dots \end{aligned}$$

In particular,

$$\begin{aligned} \Gamma(\epsilon - 1) &= -\frac{1}{\epsilon} + \gamma - 1 \\ \Gamma(\epsilon) &= \frac{1}{\epsilon} - \gamma \end{aligned}$$

Useful consequences:

$$\mu^{2\epsilon} \int \frac{d^{4-2\epsilon}q}{(2\pi)^{4-2\epsilon}} \frac{1}{q^2 + m^2} = \frac{m^2}{16\pi^2} \left[-\frac{1}{\epsilon} + \gamma - 1 - \ln 4\pi + \ln(m^2/\mu^2) \right]$$

$$\mu^{2\epsilon} \int \frac{d^{4-2\epsilon}q}{(2\pi)^{4-2\epsilon}} \frac{1}{(q^2 + m^2)^2} = \frac{1}{16\pi^2} \left[\frac{1}{\epsilon} - \gamma + \ln 4\pi - \ln(m^2/\mu^2) \right]$$